

**PHY · PHYSICS**

04. Rotational & Circular Motion

Official PMDC Solved Past Paper Topical Booklet · High-Yield Examination Series

🎯 95 Solved Questions

📅 2008 to 2024 Archive

🛡️ Boards: BUMHS, DUHS, KMU, KU, MDCAT, NUMS

📊 Weightage: 27% of MDCAT (54 MCQs)

Curated by Ibrahim for fellow bees. Every question in this chapter has been verified against official UHS, NUMS, SZABMU, DUHS, and KMU answer keys. Propolis cognitive autopsies are provided at the end of this booklet to explain core concepts and break down common distractor traps.

OFFICIAL SOLVED EDITION**Q1.**

SZABMU 2024

A rotating pulley completes twelve revolutions in 4 seconds, calculate the average angular velocity of rotating pulley in revolutions per second? [SZABMU 2024]

- (A) 3
- (B) 4
- (C) 5
- (D) 6

Q2.

SZABMU 2024

Which of the following rule helps us to detect the direction of angular velocity? [SZABMU 2024]

- (A) Head to tail rule
- (B) Kirchhoff rule
- (C) Left hand rule
- (D) Right hand rule

Q3.

SZABMU 2024

The instantaneous acceleration of an object travelling with uniform speed in a circle directed towards the center of circle is referred as ____ [SZABMU 2024]

- (A) Angular acceleration
- (B) Centrifugal acceleration
- (C) Centripetal acceleration
- (D) Tangential acceleration

Q4.

SZABMU 2024

In one dimensional elastic collision of two bodies of same masses, what will happen if moving body collides with the mass which is initially at rest? [SZABMU 2024]

- (A) The collision would become inelastic
- (B) Their velocities will be interchanged
- (C) Their velocities will remain same
- (D) Velocities of both bodies will be zero

Q5.

SZABMU 2024

Which one of the following is the SI-unit of angular displacement? [SZABMU 2024]

- (A) Degree
- (B) Radian
- (C) Revolution
- (D) Steradian

Q6.

KU 2024

A wheel makes 3 complete revolutions. What is the total number of radians through which a point on wheel has rotated? [KU 2024]

- (A) 2π
- (B) 3π
- (C) 6π
- (D) 9π

Q7.

KU 2024

In a rotating spaceship, to produce artificial gravity, what does the centripetal force do? [KU 2024]

- (A) Has no effect inside the spaceship
- (B) Increases spaceship's rotation
- (C) Pulls objects towards the centre
- (D) Pushes the objects towards the outer wall

Q8.

KU 2024

When the mass of a body moving along a circle becomes half and radius becomes double, and v is constant, the centripetal force becomes? [KU 2024]

- (A) Double
- (B) Half
- (C) One-fourth
- (D) Remains same

Q9.

KU 2024

A car is moving in a circular path at a constant speed. What provides the necessary centripetal force to keep the car moving in this path? [KU 2024]

- (A) The car's inertia resisting any change in direction
- (B) The car's mass pulling it towards the centre of the circle
- (C) The engine's power pushing the car forward
- (D) The friction between the tyres and the road

Q10.

UHS 2024

The centripetal acceleration of an object moving along a circle of radius ' r ' with an angular speed ' ω ' is given by the formula: [UHS 2024]

- (A) $a = r\omega^2$
- (B) $a = r\omega$
- (C) $a = r^2\omega$
- (D) $a = r^2\omega^2$

Q11.

UHS 2024

An air craft makes a turn in a horizontal circle of radius 100 m. It is travelling with a velocity of 250 m/s. The angular velocity of the air craft will be: [UHS 2024]

- (A) 1.5 rad/sec
- (B) 2.5 rad/sec
- (C) 3 rad/sec
- (D) 3.5 rad/sec

Q12.

UHS 2024

A particle of mass ' m ' is moving on a circular path of radius ' r ' with velocity ' v ', then centripetal force acting on it is F . If the velocity of particle increases by 2 time and radius of circular path increases by 4 times then new centripetal force F' will be: [UHS 2024]

- (A) $F' = 2F$
- (B) $F' = \frac{1}{2}F$
- (C) $F' = 4F$
- (D) $F' = F$

Q13.

UHS 2024

A roller coaster is moving with 30 ms^{-1} on a circular track of radius 30 m, then net mass of coaster + passengers is ' m ' the centripetal force acting on it is: [UHS 2024]

- (A) 900 m
- (B) m
- (C) 450 m
- (D) 30 m

Q14.

UHS 2024

In an elastic collision the total kinetic energy [UHS 2024]

- (A) Dissipates after collision
- (B) Increases after the collision
- (C) Reduces after the collision
- (D) Before and after collision remains the same

Q15.

UHS 2024

The instantaneous velocity along the curved path is: [UHS 2024]

- (A) Along the tangent
- (B) Perpendicular to the slope
- (C) Parallel to the radius
- (D) Anti-parallel to the radius

Q16.

NUMS 2024

The velocity of an object revolving in circle is: [NUMS 2024]

- (A) Constant
- (B) Variable
- (C) Zero
- (D) Negative

Q17.

NUMS 2024

Sara goes around a circular track that has diameter of 20 m. If she runs around the entire track for a distance of 80 m. What is her angular displacement? [NUMS 2024]

- (A) 16 radians
- (B) 18 radians
- (C) 50 radians
- (D) 8 radians

Q18.

NUMS 2024

Angle swept by minute hand in one minute is: [NUMS 2024]

- (A) 6°
- (B) 36°
- (C) 8°
- (D) 20°

Q19.

NUMS 2024

To find the direction of angular displacement: [NUMS 2024]

- (A) Grasp the axis of rotation in right hand
- (B) Grasp the axis of rotation in left hand
- (C) Put the thumb of right hand in the direction of circular motion
- (D) Put the thumb of left hand in the direction of circular motion

Q20.

NUMS 2024

A particle moves in a circle of radius 200 cm with a linear speed of 20 m/s. Find the angular velocity: [NUMS 2024]

- (A) 2 rad s^{-1}
- (B) 200 rad s^{-1}
- (C) 20 rad s^{-1}
- (D) 10 rad s^{-1}

Q21.

NUMS 2024

For anti-clock wise rotation the angular displacement is: [NUMS 2024]

- (A) Negative
- (B) Positive
- (C) Null
- (D) Zero

Q22.

NUMS 2024

How many degrees in $2\pi/5$ radians? [NUMS 2024]

- (A) 80°
- (B) 50°
- (C) 72°
- (D) 120°

Q23.

BUMHS 2024

Velocity 'v' of an electron revolving around the nucleus is at ____ to the radius r of the orbit. [BUMHS 2024]

- (A) Acute angle
- (B) Obtuse angle
- (C) Right angle
- (D) Supplementary angle

Q24.

BUMHS 2024

A particle is executing uniform circular motion in a circle of radius 100 mm. If its speed is 10 cm/s then its angular velocity is: [BUMHS 2024]

- (A) 10 rad/s
- (B) 0.1 rad/s
- (C) 10 revolutions/s
- (D) None of these

Q25.

BUMHS 2024

A body of mass m moving in a circle of radius is executing a uniform circular motion. If the mass of the body is doubled then the centripetal force acting upon the body is [BUMHS 2024]

- (A) Reduced to half
- (B) Remains same
- (C) Doubled
- (D) None of the given options

Q26.

UHS 2023

One radian is analogous to: [UHS 2023]

- (A) $57^\circ 3'$
- (B) $57^\circ 3''$
- (C) $57^\circ 18'$
- (D) $57^\circ 18''$

Q27.

UHS 2023

The circular line has a fix distance from ____? [UHS 2023]

- (A) Any point
- (B) A fix point
- (C) A point on a circle
- (D) A point from outside

Q28.

UHS 2023

If a particle is moving with uniform circular motion, then: [UHS 2023]

- (A) Velocity and acceleration are antiparallel
- (B) Velocity and acceleration are parallel
- (C) Velocity and acceleration are perpendicular
- (D) Zero acceleration

Q29.

UHS 2023

One degree is equal to: [UHS 2023]

- (A) $\frac{\pi}{90}$ radians
- (B) $\frac{\pi}{180}$ radians
- (C) $\frac{\pi}{270}$ radians
- (D) $\frac{\pi}{360}$ radians

Q30.

SZABMU 2023

At what angle the curved path should be banked if a car is moving with a speed of 12 m/s at a radius 26 m? [SZABMU 2023]

- (A) 10°
- (B) 15°
- (C) 20°
- (D) 22°

Q31.

SZABMU 2023

In angular motion, in one rotation there are: [SZABMU 2023]

- (A) π radians
- (B) 2π radians
- (C) $\frac{\pi}{2}$ radians
- (D) $\frac{\pi}{4}$ radians

Q32.

SZABMU 2023

If a stone of mass 1 Kg is whirled in a horizontal circle (radius = 1 m) with velocity 1 m/s, then the period is: [SZABMU 2023]

- (A) 4π
- (B) 2π
- (C) $\frac{\pi}{2}$
- (D) $\frac{2\pi}{2}$

Q33.

SZABMU 2023

The relation between linear and angular acceleration is: [SZABMU 2023]

- (A) $a = r\alpha$
- (B) $a = \frac{r}{\alpha}$
- (C) $a = r^2\alpha$
- (D) $a = r\alpha^2$

Q34.

KMU 2023

The speedometer of a car shows 36 km/hr. The angular speed of the wheels having 0.5 m radius is: [ETEA 2023]

- (A) 5 rad/sec
- (B) 10 rad/sec
- (C) 15 rad/sec
- (D) 20 rad/sec

Q35.

KMU 2023

The angular speed in radian/hour for daily rotation of our earth is: [ETEA 2023]

- (A) $\frac{\pi}{12}$
- (B) 12π
- (C) 24π
- (D) $\frac{\pi}{24}$

Q36.

KMU 2023

A fly wheel rotates at constant speed of 600 rpm. The angle described by the shaft in radian in one second is: [ETEA 2023]

- (A) 100π
- (B) 20π
- (C) 0
- (D) 20

Q37.

KMU 2023

The angular speed of the second hand of a watch is: [ETEA 2023]

- (A) $\left(\frac{\pi}{1800}\right)$ rad/sec
- (B) $\left(\frac{\pi}{60}\right)$ rad/sec
- (C) $\left(\frac{\pi}{30}\right)$ rad/sec
- (D) (2π) rad/sec

Q38.

DUHS 2023

Which of the following results in the centripetal acceleration produced in a body? [SINDH 2023]

- (A) Change in magnitude of velocity
- (B) Change in mass
- (C) Change in magnitude of force
- (D) Change in direction of velocity

Q39.

DUHS 2023

1 radian equals to: [SINDH 2023]

- (A) 0.573°
- (B) 5.73°
- (C) 57.3°
- (D) 573°

Q40.

DUHS 2023

A wheel of bike rotates from rest and achieves 6.28 radian/sec in 2 sec. What will be its angular acceleration? [SINDH 2023]

- (A) 1 rad/sec²
- (B) 1.5 rad/sec²
- (C) 3.14 rad/sec²
- (D) 6.28 rad/sec²

Q41.

DUHS 2023

The ratio between linear velocity and radius equals to: [SINDH 2023]

- (A) Angular displacement
- (B) Angular velocity
- (C) Angular acceleration
- (D) Angular momentum

Q42.

NUMS 2023

The relation between radian and degree is: [NUMS 2023]

- (A) 1 rad = 57.3°
- (B) $1^\circ = 57.3$ rad
- (C) 1 rad = 1°
- (D) $1^\circ = \pi$ rad

Q43.

NUMS 2023

In case of centripetal force the value of instantaneous acceleration is given by: [NUMS 2023]

- (A) $a_c = \frac{v}{r}$
- (B) $a_c = \frac{v^2}{r}$
- (C) $a_c = vr$
- (D) $a_c = v^2 r$

Q44.

NUMS 2023

An electric motor turns at 400 revolutions per minute. Its angular velocity in rad/s will be: [NUMS 2023]

- (A) $\frac{20\pi}{3}$
- (B) $\frac{30\pi}{3}$
- (C) $\frac{40\pi}{3}$
- (D) $\frac{30\pi}{4}$

Q45.

NUMS 2023

The expression for centripetal force is: [NUMS 2023]

- (A) $F_c = m\omega^2$
- (B) $F_c = mr\omega$
- (C) $F_c = mr\omega^2$
- (D) $F_c = mr^2\omega$

Q46.

UHS 2022

A fireman wants to slide down a rope. The breaking load of the rope is $\frac{3}{4}$ th of the weight of the man. With what acceleration should the fire man slide down? (Acceleration due to gravity is 'g') [UHS 2022]

- (A) g
- (B) $\frac{g}{4}$
- (C) $\frac{3g}{4}$
- (D) 0

Q47.

UHS 2022

The number of revolutions in 3π radians is [UHS 2022]

- (A) $\frac{1}{60}$
- (B) $\frac{3}{2}$
- (C) 2
- (D) 6

Q48.

UHS 2022

If a flywheel is rotating at 3.0 rad/s, the time it takes to complete one revolution is about [UHS 2022]

- (A) 0.67 s
- (B) 1.3 s
- (C) 1.0 s
- (D) 2.1 s

Q49.

UHS 2022

A fighter plane is moving in a vertical circle of radius r. Its minimum velocity at the highest point of the circle will be? [UHS 2022]

- (A) $\sqrt{3gr}$
- (B) $\sqrt{2gr}$
- (C) $\sqrt{gr}$
- (D) $\sqrt{\frac{gr}{2}}$

Q50.

SZABMU 2022

Which one of the following provides centripetal force in a circular motion of body? [SZABMU 2022]

- (A) Vertical component of weight
- (B) Horizontal component of weight
- (C) Weight of the body
- (D) Force of friction

Q51.

SZABMU 2022

How many radians are in one degree? [SZABMU 2022]

- (A) 0.0174 rad
- (B) 0.174 rad
- (C) 1.745 rad
- (D) 0.00174 rad

Q52.

SZABMU 2022

Which one of the following is the angular velocity of an electric motor if it runs at 400 rpm? [SZABMU 2022]

- (A) 51.2 rads^{-1}
- (B) 41.9 rads^{-1}
- (C) 45.2 rads^{-1}
- (D) 38.5 rads^{-1}

Q53.

SZABMU 2022

The tension in string at top of vertical circle is: [SZABMU 2022]

- (A) Zero
- (B) mg
- (C) $2mg$
- (D) $3mg$

Q54.

SZABMU 2022

The centripetal force in terms of angular velocity is given by [SZABMU 2022]

- (A) $F_c = mr\omega$
- (B) $F_c = mr\omega^2$
- (C) $F_c = mr\alpha^2$
- (D) $F_c = mr\alpha$

Q55.

SZABMU 2022

Which one of the following is angular speed in radian per hour for daily rotation of our earth? [SZABMU 2022]

- (A) 2π
- (B) 4π
- (C) $\frac{\pi}{6}$
- (D) $\frac{\pi}{12}$

Q56.

KMU 2022

Which one is not a vector quantity? [ETEA 2022]

- (A) Angular displacement
- (B) Impulse
- (C) Moment of inertia
- (D) Momentum

Q57.

KMU 2022

When angular speed of a body is doubled, the centripetal acceleration becomes: [ETEA 2022]

- (A) Doubled
- (B) Thrice
- (C) Four times
- (D) Remains the same

Q58.

BUMHS 2022

According to the right-hand rule, the direction of angular displacement is along the? [BMU 2022]

- (A) Curl of fingers
- (B) First finger
- (C) Pointing thumb
- (D) Second finger

Q59.

BUMHS 2022

The ratio of angular speed of a minute hand to hour hand of the watch is? [BMU 2022]

- (A) 1:6
- (B) 6:1
- (C) 1:12
- (D) 12:1

Q60.

BUMHS 2022

In the SI system of units, the centripetal force is measured in: [BMU 2022]

- (A) J
- (B) N
- (C) T
- (D) W

Q61.

BUMHS 2022

The best example of uniform circular motion is: [BMU 2022]

- (A) The motion of the earth around the sun
- (B) The motion of a mass attached with a string on a vertical circle
- (C) The motion of a minute hand in the clock
- (D) The motion of a spinning top

Q62.

DUHS 2022

A cyclist moving on a circular track skid because:
[DUHS 2022]

- (A) The limiting friction is perpendicular to the centripetal force
- (B) The limiting friction is greater than the required centripetal force
- (C) The limiting friction is opposite to the centripetal force
- (D) The limiting friction is less than the required centripetal force

Q63.

DUHS 2022

Every particle of a rotating disc moves with: [DUHS 2022]

- (A) Constant linear velocity
- (B) Constant angular velocity
- (C) Constant tangential velocity
- (D) Constant linear displacement

Q64.

DUHS 2022

A body is moving in a circle with constant speed, with radius of circle is 'r' and time period 'T' the centripetal acceleration is given by: [DUHS 2022]

- (A) $a_c = \frac{4\pi r^2}{T^2}$
- (B) $a_c = \frac{4\pi^2 r}{T^2}$
- (C) $a_c = \frac{4\pi r}{T}$
- (D) $a_c = \frac{4\pi^2 r}{T}$

Q65.

DUHS 2022

The angular momentum of a particle changes from 0 to 720 J s in 4 s. The magnitude of torque is: [DUHS 2022]

- (A) 2880 N-m
- (B) 180 N-m
- (C) 360 N-m
- (D) 1440 N-m

Q66.

NUMS 2022

1 radian = [NUMS 2022]

- (A) 53.7°
- (B) 73.5°
- (C) 57.3°
- (D) 37.5°

Q67.

NUMS 2022

The minimum required velocity to put a satellite into the orbit is called: [NUMS 2022]

- (A) Terminal velocity
- (B) Escape velocity
- (C) Critical velocity
- (D) Average velocity

Q68.

NUMS 2022

A rotating wheel complete 12 rev in 4s. Find the average angular velocity its rad/sec. [NUMS 2022]

- (A) 24.6 rad/s
- (B) 18.5 rad/s
- (C) 18.8 rad/s
- (D) 10.4 rad/s

Q69.

NUMS 2022

When an object experiences circular motion, the directional of centripetal acceleration is: [NUMS 2022]

- (A) Toward center
- (B) Along tangent
- (C) Along the direction of motion
- (D) Opposite the direction of motion

Q70.

PMDC 2021

Number of revolutions in 720 degrees. [NMDCAT 2021]

- (A) 4 revolutions
- (B) 3 revolutions
- (C) 2 revolutions
- (D) 9 revolutions

Q71.

PMDC 2021

Car moves in circular path due to [NMDCAT 2021]

- (A) Centripetal force
- (B) Frictional force
- (C) Gravitational force
- (D) None

Q72.

PMDC 2021

Radian is used to measure [NMDCAT 2021]

- (A) Small angles
- (B) Large angles
- (C) Both
- (D) None

Q73.

PMDC 2021

If earth covers 360 degrees in 24 hours, then how many degrees are in two hours [NMDCAT 2021]

- (A) 15°
- (B) 30°
- (C) 45°
- (D) 60°

Q74.

PMDC 2021

Earth completes one revolution about its own axis [NMDCAT 2021]

- (A) 24 hours
- (B) 6 months
- (C) 12 months
- (D) None

Q75.

PMDC 2020

If during circular motion, tangential velocity of a body becomes double, then centripetal force becomes: [NMDCAT 2020]

- (A) Double
- (B) One half
- (C) Four times
- (D) One fourth

Q76.

PMDC 2020

Under what condition an object will have zero displacement but non zero distance? [NMDCAT 2020]

- (A) Linear motion
- (B) Circular motion
- (C) Random motion
- (D) Oscillation

Q77.

PMDC 2020

1 radian is equal to [NMDCAT 2020]

- (A) 57.1°
- (B) 57.2°
- (C) 57.3°
- (D) 57.4°

Q78.

NUMS 2020

A 10 N force moves a body around a circular path of radius 50 cm. What is work done in completing one revolution? [NUMS 2020]

- (A) 5 J
- (B) 0
- (C) 31.42 J
- (D) 500 J

Q79.

MDCAT 2019

An object is moving along a circular path of radius 4 m. What will be its angular displacement if it moves 14 m on this circular path? [MDCAT 2019]

- (A) 5.5 radians
- (B) 5.0 radians
- (C) 3.5 radians
- (D) 4.5 radians

Q80.

MDCAT 2018

A wheel starts rotating from rest with angular acceleration of 2 rad s^{-2} till its angular speed becomes 6 rad/s. The angular displacement of the wheel will be equal to [MDCAT 2018]

- (A) 4 rad
- (B) 12 rad
- (C) 9 rad
- (D) 7 rad

Q81.

MDCAT 2018

Which of the following gives the relationship between linear velocity and angular velocity? [MDCAT 2018]

- (A) $v = r\omega$
- (B) $v = s\omega$
- (C) $v = r\theta$
- (D) $v = s\theta$

Q82.

MDCAT 2017

A body moves in a circle with increasing angular velocity. At time $t = 6 \text{ sec}$, the angular velocity is 27 rad/s. What is the radius of circle made by the body where linear velocity is 81 cm/s? [MDCAT 2017]

- (A) 6 cm
- (B) 9 cm
- (C) 3 cm
- (D) 7 cm

Q83.

MDCAT 2017

Angular speed of minutes hand of mechanical watch is: [MDCAT 2017]

- (A) $\frac{\pi}{30} \text{ rad min}^{-1}$
- (B) $\frac{\pi}{2} \text{ rad min}^{-1}$
- (C) $\pi \text{ rad min}^{-1}$
- (D) None of these

Q84.

MDCAT 2017

A body is moving in a circular path with constant speed. The magnitude of tangential and centripetal acceleration are: **[MDCAT 2017]**

- (A) Tangential: rv^2 Centripetal: 0
- (B) Tangential: 0 Centripetal: $\frac{v^2}{r}$
- (C) Tangential: 0 Centripetal: 0
- (D) Tangential: $\frac{v^2}{r}$ Centripetal: $\frac{v^2}{r}$

Q85.

KMU 2017

A centripetal force F acts on a body moving with angular speed ω . If the angular speed is tripled, then the magnitude of centripetal force becomes: **[ETEA 2017]**

- (A) $8F$
- (B) $9F$
- (C) $3F$
- (D) $4F$

Q86.

KMU 2017

A fly wheels rotates at a constant speed of 3000 rpm (rev/min). The angle described by the shaft in radian in one second is: **[ETEA 2017]**

- (A) 2π
- (B) 30π
- (C) 100π
- (D) 300π

Q87.

KMU 2016

A wheel starts from rest and has an angular acceleration of 4.0 rad/s^2 . When it has made 10 rev its angular velocity is: **[ETEA 2016]**

- (A) 16 rad/s
- (B) 22 rad/s
- (C) 32 rad/s
- (D) 250 rad/s

Q88.

KMU 2016

A child riding on a large merry-go-round, travels a distance of 3000 m in a circle of diameter 40 m. The total angle through which she revolves is: **[ETEA 2016]**

- (A) 50 rad
- (B) 75 rad
- (C) 150 rad
- (D) 314 rad

Q89.

KMU 2016

The angular velocity for daily rotation of the earth is: **[ETEA 2016]**

- (A) $\frac{\pi}{3} \text{ radian hr}^{-1}$
- (B) $\frac{\pi}{6} \text{ radian hr}^{-1}$
- (C) $\frac{\pi}{12} \text{ radian hr}^{-1}$
- (D) $12\pi \text{ radian hr}^{-1}$

Q90.

KMU 2013

The centripetal acceleration of a car travelling at constant speed around a frictionless circular track: **[ETEA 2013]**

- (A) Is zero
- (B) Has constant magnitude but varying direction
- (C) Has constant direction but varying magnitude
- (D) Has varying magnitude and direction

Q91.

MDCAT 2011

Radian is a unit of angular displacement which can also be measured in degrees. How many radians are equal to one degree? **[MCAT 2011]**

- (A) $\frac{180}{\pi}$
- (B) $\frac{\pi}{180}$
- (C) $\frac{2\pi}{180}$
- (D) $\frac{\pi}{57.3}$

Q92.

MDCAT 2010

A wheel of radius 1 m covers an angular displacement of 180° . Its linear displacement is: **[MCAT 2010]**

- (A) 3.14 m
- (B) 6.28 m
- (C) $\pi \text{ rad}$
- (D) 0.157 m

Q93.

KMU 2010

When a body moves in a circle the angle between its linear velocity and angular velocity is always: **[ETEA 2010]**

- (A) 0°
- (B) 180°
- (C) 360°
- (D) 90°

Q94.

MDCAT 2009

Linear velocity or tangential velocity of any particle moving in a circular path of radius 2 m with angular velocity 8 rad s^{-1} will be: [MCAT 2009]

- (A) 16 ms^{-1}
- (B) 10 ms^{-1}
- (C) 4 ms^{-1}
- (D) 6 ms^{-1}

Q95.

MDCAT 2008

The force required to bend the normally straight path of a particle into a circular path is called ____ force. [MCAT 2008]

- (A) Traveling
- (B) Centrifugal
- (C) Bending
- (D) Centripetal

RAPID 5–COLUMN ANSWER KEY

Official PMDC verified answer keys for 04. Rotational & Circular Motion.

beambeprep.com

Q#	KEY	BOARD	Q#	KEY	BOARD	Q#	KEY	BOARD	Q#	KEY	BOARD	Q#	KEY	BOARD
1	A	SZABMU 2024	20	D	NUMS 2024	39	C	DUHS 2023	58	C	BUMHS 2022	77	C	PMDC 2020
2	D	SZABMU 2024	21	B	NUMS 2024	40	C	DUHS 2023	59	D	BUMHS 2022	78	B	NUMS 2020
3	C	SZABMU 2024	22	C	NUMS 2024	41	B	DUHS 2023	60	B	BUMHS 2022	79	C	MDCAT 2019
4	B	SZABMU 2024	23	C	BUMHS 2024	42	A	NUMS 2023	61	C	BUMHS 2022	80	C	MDCAT 2018
5	B	SZABMU 2024	24	D	BUMHS 2024	43	B	NUMS 2023	62	D	DUHS 2022	81	A	MDCAT 2018
6	C	KU 2024	25	C	BUMHS 2024	44	C	NUMS 2023	63	B	DUHS 2022	82	C	MDCAT 2017
7	C	KU 2024	26	C	UHS 2023	45	C	NUMS 2023	64	B	DUHS 2022	83	A	MDCAT 2017
8	C	KU 2024	27	B	UHS 2023	46	B	UHS 2022	65	B	DUHS 2022	84	B	MDCAT 2017
9	D	KU 2024	28	C	UHS 2023	47	B	UHS 2022	66	C	NUMS 2022	85	B	KMU 2017
10	A	UHS 2024	29	B	UHS 2023	48	D	UHS 2022	67	C	NUMS 2022	86	C	KMU 2017
11	B	UHS 2024	30	D	SZABMU 2023	49	C	UHS 2022	68	C	NUMS 2022	87	B	KMU 2016
12	D	UHS 2024	31	B	SZABMU 2023	50	D	SZABMU 2022	69	A	NUMS 2022	88	C	KMU 2016
13	D	UHS 2024	32	B	SZABMU 2023	51	A	SZABMU 2022	70	C	PMDC 2021	89	C	KMU 2016
14	D	UHS 2024	33	A	SZABMU 2023	52	B	SZABMU 2022	71	B	PMDC 2021	90	B	KMU 2013
15	A	UHS 2024	34	D	KMU 2023	53	A	SZABMU 2022	72	B	PMDC 2021	91	B	MDCAT 2011
16	B	NUMS 2024	35	A	KMU 2023	54	B	SZABMU 2022	73	B	PMDC 2021	92	A	MDCAT 2010
17	D	NUMS 2024	36	B	KMU 2023	55	D	SZABMU 2022	74	A	PMDC 2021	93	D	KMU 2010
18	A	NUMS 2024	37	C	KMU 2023	56	C	KMU 2022	75	C	PMDC 2020	94	A	MDCAT 2009
19	A	NUMS 2024	38	D	DUHS 2023	57	C	KMU 2022	76	B	PMDC 2020	95	D	MDCAT 2008

TOTAL
MCQS

95

CORRECT

INCORRECT

BLANK

ACCURACY

%

Scan the booklet QR code to enter answers in
BeambePrep Grind Mode for automated error
 autopsies.

PROPOLIS COGNITIVE ERROR AUTOPSIES

Step-by-step conceptual derivations, formula working, and distractor autopsies for every question.

Question #1 Key: (A)

SZABMU 2024

A rotating pulley completes twelve revolutions in [Formula], calculate the average angular velocity of rotation.

CORE CONCEPT AND DERIVATION

Angular velocity can be expressed in terms of frequency (revolutions per second), which is simply the total revolutions divided by total time.

Formula:

$$f = \frac{\text{Total Revolutions}}{\text{Total Time}}$$

Solution:

- Given total revolutions = 12.
- Given total time = 4 seconds.
- Divide the two: $f = \frac{12}{4} = 3 \text{ rev/sec.}$

Why other options are incorrect:

- **Option B, Option C, Option D** are mathematically incorrect calculations for $12 \div 4$.

Question #2 Key: (D)

SZABMU 2024

Which of the following rule helps us to detect the direction of angular velocity? [SZABMU 2024]

CORE CONCEPT AND DERIVATION

Rotational vectors (like torque, angular momentum, and angular velocity) are 'pseudo-vectors'. Their direction points along the axis of rotation, defined by a universal human convention.

Solution:

- By universal physics convention, the **Right Hand Rule** is used.
- If you curl the fingers of your right hand in the direction the object is physically spinning, your extended thumb points strictly along the axis in the direction of the angular velocity vector.

Why other options are incorrect:

- **Head to tail rule** is used for adding linear geometric vectors.
- **Kirchhoff's rule** is for electrical circuits.
- **Left hand rule** is primarily used for magnetic force on a moving charge (Fleming's rule).

Question #3 Key: (C)

SZABMU 2024

The instantaneous acceleration of an object travelling with uniform speed in a circle directed towards the center.

CORE CONCEPT AND DERIVATION

In uniform circular motion, an object is continually turning inward to stay on the circle, implying a continuous inward acceleration vector.

Solution:

- The term 'centripetal' stems from Latin words literally translating to 'center-seeking'.
- Thus, the specific acceleration vector strictly pointing towards the center of curvature is universally called **Centripetal acceleration**.

Why other options are incorrect:

- **Option A** relates to the change in rotational speed (spin rate), which is zero here.
- **Option B** 'Centrifugal' means center-fleeing (outward), which is a fictitious pseudo-force/acceleration in a rotating frame.
- **Option D** relates to speeding up or slowing down along the track.

Question #4 Key: (B)

SZABMU 2024

In one dimensional elastic collision of two bodies of same masses, what will happen if moving body collides wi...

CORE CONCEPT AND DERIVATION

In a perfectly elastic, 1-dimensional collision between two perfectly identical masses, both momentum and kinetic energy are conserved.

Solution:

- When we solve the simultaneous conservation equations (Momentum: $mv_1 = mv'_1 + mv'_2$ and KE: $\frac{1}{2}mv_1^2 = \frac{1}{2}mv_1'^2 + \frac{1}{2}mv_2'^2$), a unique mathematical symmetry emerges.
- The solution mandates that the objects perfectly swap their initial velocity states.
- The moving body stops completely, transferring 100% of its momentum to the second body, which shoots forward with the original velocity.
- Thus, their velocities are exactly **interchanged**.

Why other options are incorrect:

- **Option A** violates the prompt which states it is elastic.
- **Option C** means the moving body passed through the stationary one like a ghost.
- **Option D** violates conservation of momentum.

Question #5 Key: (B)

SZABMU 2024

Which one of the following is the SI-unit of angular displacement? [SZABMU 2024]

CORE CONCEPT AND DERIVATION

Angular displacement requires a standardized dimensionless unit in the International System (SI) to easily interface with linear kinematics formulas without messy conversion constants.

Solution:

- The **Radian** is defined as the ratio of arc length to radius (S/r). Since it is a ratio of lengths (meters/meters), it is a pure, dimensionless number, making it the mathematically elegant and official SI unit for planar angles.

Why other options are incorrect:

- **Option A** and **Option C** are common practical units but are not the official SI standard.
- **Option D** (Steradian) is the SI unit for *solid* (3D) angles, not standard 2D angular displacement.

Question #6 Key: (C)

KU 2024

A wheel makes 3 complete revolutions. What is the total number of radians through which a point on wheel has r...

CORE CONCEPT AND DERIVATION

One full revolution around a circle corresponds to the complete circumference arc length, which translates strictly to 2π radians.

Formula:

$$\theta_{\text{total}} = N \times 2\pi$$

Solution:

- Number of revolutions $N = 3$.
- Multiply by the radian value of one revolution: $3 \times 2\pi = 6\pi$ radians.

Why other options are incorrect:

- **Option A** represents just 1 revolution.
- **Option B** represents 1.5 revolutions.
- **Option D** represents 4.5 revolutions.

Question #7 Key: (C)

KU 2024

In a rotating spaceship, to produce artificial gravity, what does the centripetal force do? [KU 2024]

CORE CONCEPT AND DERIVATION

Artificial gravity utilizes inertia. The spaceship spins, meaning everything inside wants to fly off in a straight straight line. The outer hull of the ship gets in the way, pushing inward on the astronauts.

Solution:

- The normal force exerted by the rigid floor/wall of the spaceship acts universally inward, serving as the required **centripetal force**.
- It physically **pulls/pushes objects strictly towards the center** of rotation to maintain their circular path.
- (Note: The astronauts' inertia pushing *outward* against the floor is the reactive centrifugal effect they feel as "gravity", but the centripetal force itself points inward).

Why other options are incorrect:

- **Option D** describes the apparent centrifugal pseudo-force experienced in the rotating frame, not the centripetal force.
- **Option B** violates conservation of angular momentum.

Question #8 Key: (C)

KU 2024

When the mass of a body moving along a circle becomes half and radius becomes double, and v is constant, the $c...$

CORE CONCEPT AND DERIVATION

Centripetal force depends directly on mass and inversely on the radius. We track these changes by plugging the new variables into the formula.

Formula:

$$F_c = \frac{mv^2}{r}$$

Solution:

- Let the initial force be $F = \frac{mv^2}{r}$.
- The new mass is $\frac{m}{2}$ and the new radius is $2r$.
- Substitute these into the formula: $F' = \frac{(\frac{m}{2})v^2}{2r}$.
- Simplify the fraction: $F' = \frac{mv^2}{2 \times 2r} = \frac{mv^2}{4r}$.
- Extract the original formula: $F' = \frac{1}{4} \left(\frac{mv^2}{r} \right) = \frac{1}{4} F$.
- Therefore, the force becomes one-fourth.

Why other options are incorrect:

- Option D** happens if you mistakenly think halving mass and doubling radius cancel each other out (they compound, not cancel, since radius is in the denominator).

Question #9 Key: (D)

KU 2024

A car is moving in a circular path at a constant speed. What provides the necessary centripetal force to keep ...

CORE CONCEPT AND DERIVATION

Centripetal force must be provided by a real, physical interaction. Without a physical force pulling sideways, an object will move in a straight line.

Solution:

- When a car turns its wheels, the rubber of the tires attempts to slide sideways across the asphalt.
- The microscopic grip of the road opposes this slide.
- This grip is the **static friction between the tires and the road**, and it pushes horizontally inward toward the center of the curve, providing the required centripetal force.

Why other options are incorrect:

- Option A** tries to go straight, which actively opposes circular motion.
- Option B** describes gravity, which pulls straight down, not sideways.
- Option C** provides forward tangential velocity, not inward turning force.

Question #10 Key: (A)

UHS 2024

The centripetal acceleration of an object moving along a circle of radius ' r ' with an angular speed ' ω ' [Formula...]

CORE CONCEPT AND DERIVATION

Centripetal acceleration is standardly derived by substituting the relationship between linear and angular speed into the primary kinematic formula.

Formula:

$$a_c = \frac{v^2}{r} \quad \text{and} \quad v = r\omega$$

Solution:

- Substitute $(r\omega)$ into the acceleration formula: $a_c = \frac{(r\omega)^2}{r}$.
- Square the terms in the numerator: $a_c = \frac{r^2\omega^2}{r}$.
- Cancel one ' r ': $a_c = r\omega^2$.

Why other options are incorrect:

- Option B** defines linear velocity, not acceleration.
- Option C** and **Option D** feature an incorrectly squared radius term.

Question #11 Key: (B)

UHS 2024

An air craft makes a turn in a horizontal circle of radius [Formula]. It is travelling with a velocity of [Fo...]

CORE CONCEPT AND DERIVATION

Angular velocity represents how fast an object is sweeping through angles around the center. It is a direct ratio of linear speed to the path's radius.

Formula:

$$\omega = \frac{v}{r}$$

Solution:

- Given linear velocity $v = 250$ m/s.
- Given radius $r = 100$ m.
- Substitute into the formula: $\omega = \frac{250}{100}$.
- $\omega = 2.5$ rad/sec.

Why other options are incorrect:

- Option A**, **Option C**, **Option D** are mathematically incorrect calculations for the simple division of 250 by 100.

Question #12 Key: (D)

UHS 2024

A particle of mass 'm' is moving on a circular path of radius 'r' with velocity 'v', then centripetal force ac...

CORE CONCEPT AND DERIVATION

By updating the variables in the centripetal force equation, we can see how simultaneous changes offset or amplify each other.

Formula:

$$F = \frac{mv^2}{r}$$

Solution:

- Let the new velocity be $v' = 2v$.
- Let the new radius be $r' = 4r$.
- Substitute these into the force equation: $F' = \frac{m(2v)^2}{4r}$.
- Square the velocity term: $F' = \frac{m(4v^2)}{4r}$.
- The 4 in the numerator and the 4 in the denominator perfectly cancel out: $F' = \frac{mv^2}{r}$.
- This is strictly equal to the original force F .

Why other options are incorrect:

- **Option A, Option B, Option C** fail to correctly square the velocity change or miss the cancellation algebra.

Question #13 Key: (D)

UHS 2024

A roller coaster is moving with [Formula] on a circular track of radius [Formula], then net mass of coaster +...

CORE CONCEPT AND DERIVATION

We calculate centripetal force by keeping the unknown mass variable 'm' as an algebraic coefficient in our final answer.

Formula:

$$F_c = \frac{mv^2}{r}$$

Solution:

- Given velocity $v = 30$ m/s.
- Given radius $r = 30$ m.
- Substitute into the formula: $F_c = \frac{m(30)^2}{30}$.
- The square on the top cancels out the 30 on the bottom.
- $F_c = m(30)$, which is typically written algebraically as $30m$.

Why other options are incorrect:

- **Option A** occurs if you square the 30 but forget to divide by the radius.
- **Option B** occurs if you assume velocity and radius cancel completely without squaring.

Question #14 Key: (D)

UHS 2024

In an elastic collision the total kinetic energy [UHS 2024]

CORE CONCEPT AND DERIVATION

Collisions are categorized by how they handle energy. In all collisions, momentum is conserved. But Kinetic Energy is treated differently.

Solution:

- By strict physical definition, a perfectly **elastic** collision is one where 0% of the kinetic energy is converted to heat, sound, or deformation.
- Therefore, the total kinetic energy of the system strictly **remains the same** before and after the collision.

Why other options are incorrect:

- **Option A** and **Option C** describe an *inelastic* collision (like cars crashing and crumpling).
- **Option B** would violate the First Law of Thermodynamics (energy cannot be created) unless there was an explosion.

Question #15 Key: (A)

UHS 2024

The instantaneous velocity along the curved path is: [UHS 2024]

CORE CONCEPT AND DERIVATION

Velocity is a vector pointing in the exact direction an object is heading at any given frozen split-second in time.

Solution:

- If the centripetal force suddenly vanished (e.g., a string snaps), inertia dictates the object will fly off in a straight line.
- This straight line only touches the curve at exactly one mathematical point.
- A line touching a circle at exactly one point is called a **tangent**.
- Thus, instantaneous velocity always points along the tangent.

Why other options are incorrect:

- **Option B** is physically meaningless terminology in this context.
- **Option C** and **Option D** describe radial motion (falling in or flying directly outward from the center), completely ignoring the forward motion.

Question #16 Key: (B)

NUMS 2024

The velocity of an object revolving in circle is: [NUMS 2024]

CORE CONCEPT AND DERIVATION

Velocity is a vector quantity, meaning it is defined by two strictly vital components: magnitude (speed) and direction.

Solution:

- Even if the object's speed (magnitude) remains perfectly constant, its path is circular.
- A circular path means the object is constantly changing the spatial direction it is pointing.
- Since the direction continuously changes, the velocity vector continuously changes.
- Therefore, velocity is considered **variable**.

Why other options are incorrect:

- **Option A** incorrectly conflates scalar 'speed' with vector 'velocity'.
- **Option C** implies the object isn't moving.
- **Option D** is meaningless since vectors change between positive and negative depending on the quadrant.

Question #17 Key: (D)

NUMS 2024

Sara goes around a circular track that has diameter of [Formula] . If she runs around the entire track for a d...

CORE CONCEPT AND DERIVATION

Angular displacement evaluates the total distance traversed strictly against the radius of the circle.

Formula:

$$\theta = \frac{S}{r}$$

Solution:

- The diameter is 20 m, so the radius $r = \frac{20}{2} = 10$ m.
- The linear distance traversed $S = 80$ m.
- Substitute into the formula: $\theta = \frac{80}{10}$.
- This gives exactly 8 radians.

Why other options are incorrect:

- **Option A** occurs if you wrongly divide the distance by the diameter ($80/20 = 4$) and then randomly multiply by something, or misapply formulas. Wait, actually, if you use diameter, you get 4. No clear logic leads to A, B, or C except random guessing.

Question #18 Key: (A)

NUMS 2024

Angle swept by minute hand in one minute is: [NUMS 2024]

CORE CONCEPT AND DERIVATION

We determine rate of motion for clock hands by dividing a full circle by the total time it takes the hand to complete that circle.

Formula:

$$\text{Rate} = \frac{360^\circ}{T}$$

Solution:

- A minute hand takes exactly 60 minutes to complete one full revolution around the clock face.
- A full revolution is 360° .
- Divide total angle by total time: $\frac{360^\circ}{60 \text{ min}} = 6^\circ$ per minute.

Why other options are incorrect:

- **Option B** would mean it finishes the circle in 10 minutes.
- **Option C** and **Option D** do not mathematically align with base-60 clock divisions.

Question #19 Key: (A)

NUMS 2024

To find the direction of angular displacement: [NUMS 2024]

CORE CONCEPT AND DERIVATION

Physics relies on standard conventions to assign a unified direction to rotational vectors (angular displacement, velocity, acceleration).

Solution:

- This convention is strictly the **Right Hand Rule**.
- You must **grasp the axis of rotation in your right hand** so that your curled fingers mimic the direction of the physical spin.
- Your extended thumb will uniquely point in the direction of the angular vector.

Why other options are incorrect:

- **Option B** and **Option D** use the left hand, which would yield the exact opposite (anti-parallel) vector.
- **Option C** incorrectly suggests placing the thumb along the curved path of motion, whereas the fingers belong on the path and the thumb belongs straight on the axis.

Question #20 Key: (D)

NUMS 2024

A particle moves in a circle of radius [Formula] with a linear speed of [Formula]. Find the angular velocity:...

CORE CONCEPT AND DERIVATION

Angular velocity is linearly proportional to speed but inversely proportional to radius. Units must be carefully harmonized into standard SI (meters) before calculating.

Formula:

$$\omega = \frac{v}{r}$$

Solution:

- First, convert the radius into meters: 200 cm = 2 meters.
- Given linear speed $v = 20$ m/s.
- Substitute into the formula: $\omega = \frac{20}{2}$.
- $\omega = 10$ rad s⁻¹.

Why other options are incorrect:

- **Option B** occurs if you forget to convert centimeters to meters and perform $\frac{20}{200}$ backward, or some other math error.
- **Option C** just repeats the linear speed.
- **Option A** just repeats the converted radius.

Question #21 Key: (B)

NUMS 2024

For anti-clock wise rotation the angular displacement is: [NUMS 2024]

CORE CONCEPT AND DERIVATION

By applying the right-hand rule to a 2D Cartesian plane (x-y axis), we establish the universal mathematical sign convention for rotation.

Solution:

- If an object rotates anti-clockwise (counter-clockwise) on a piece of paper, your right fingers curl with the motion, and your thumb points directly UP toward your face (positive z-axis).
- Therefore, in mathematics and physics, anti-clockwise rotation is universally defined as **Positive**.
- Clockwise rotation points the thumb down into the paper and is defined as Negative.

Why other options are incorrect:

- **Option A** represents clockwise rotation.
- **Option C** and **Option D** imply the object has not rotated at all.

Question #22 Key: (C)

NUMS 2024

How many degrees in [Formula] ? [NUMS 2024]

CORE CONCEPT AND DERIVATION

To convert radians to degrees, we can safely replace the constant π with its degree-equivalent of a half-circle (180°).

Formula:

$$\text{Degrees} = \text{Radians} \times \frac{180^\circ}{\pi}$$

Solution:

- Given angle: $\frac{2\pi}{5}$.
- Substitute $\pi = 180^\circ$.
- Calculation: $\frac{2 \times 180^\circ}{5}$.
- $\frac{360^\circ}{5} = 72^\circ$.

Why other options are incorrect:

- **Option D** is $2\pi/3$ radians.
- **Option A** and **Option B** are mathematical errors when dividing 360 by 5.

Question #23 **Key: (C)**

BUMHS 2024

Velocity 'v' of an electron revolving around the nucleus is at ____ to the radius r of the orbit. [BUMHS 2024]

CORE CONCEPT AND DERIVATION

This applies basic circular kinematics to a planetary/Bohr atomic model. The direction of instantaneous linear velocity relative to the center governs the shape of the orbit.

Solution:

- In uniform circular motion, the radius line points from the center strictly outward.
- The velocity vector points strictly along the tangent line to the circular path.
- By geometric theorems, a tangent line to a circle is always perfectly perpendicular to the radius line connecting the tangent point to the center.
- Therefore, the angle between them is exactly 90° , which is a **Right angle**.

Why other options are incorrect:

- **Option A** or **Option B** would imply the electron is spiraling inward or spiraling outward, escaping the orbit.
- **Option D** refers to two angles summing to 180, which doesn't directly describe a single geometric vector intersection.

Question #24 **Key: (D)**

BUMHS 2024

A particle is executing uniform circular motion in a circle of radius [Formula]. If its speed is [Formula] th...

CORE CONCEPT AND DERIVATION

To calculate angular velocity, all spatial dimensions must be equalized. We must convert millimeters and centimeters to a common unit.

Formula:

$$\omega = \frac{v}{r}$$

Solution:

- Let's convert both into centimeters.
- Radius $r = 100 \text{ mm} = 10 \text{ cm}$.
- Speed $v = 10 \text{ cm/s}$.
- Substitute into formula: $\omega = \frac{10 \text{ cm/s}}{10 \text{ cm}}$.
- $\omega = 1 \text{ rad/s}$.
- Since 1 rad/s is not present in Options A, B, or C, the correct choice must be "None of these".

Why other options are incorrect:

- **Option A** happens if you wrongly divide 100 by 10.
- **Option B** happens if you wrongly divide 10 by 100 (forgetting to convert mm to cm).
- **Option C** mistakes radians for full revolutions, which are off by a factor of 2π .

Question #25 **Key: (C)**

BUMHS 2024

A body of mass m moving in a circle of radius r is executing a uniform circular motion. If the mass of the body ...

CORE CONCEPT AND DERIVATION

The centripetal force required to hold an object in a circle is directly, linearly proportional to the mass of the object, assuming speed and radius are unchanged.

Formula:

$$F_c = \frac{mv^2}{r}$$

Solution:

- Let initial force be $F = \frac{mv^2}{r}$.
- If the new mass is $2m$, substitute it into the formula.
- New force $F' = \frac{(2m)v^2}{r} = 2 \left(\frac{mv^2}{r} \right)$.
- Therefore, $F' = 2F$, meaning the required force is **Doubled**.

Why other options are incorrect:

- Option A** assumes an inverse relationship.
- Option B** assumes force is independent of mass (which is only true for the *acceleration* of an object falling under gravity, but not the physical force required to bend its path).

Question #26 **Key: (C)**

UHS 2023

One radian is analogous to: [UHS 2023]

CORE CONCEPT AND DERIVATION

A radian is the standard unit of angular measure. To convert radians to degrees, minutes, and seconds, we multiply by the conversion factor $180/\pi$.

Formula:

$$1 \text{ rad} = \frac{180^\circ}{\pi}$$

Solution:

- We know $1 \text{ rad} \approx 57.2958^\circ$.
- This can be split into exactly 57° and a fractional remainder of 0.2958° .
- To find the minutes, multiply the fraction by 60: $0.2958 \times 60 \approx 17.75'$.
- Rounding to the nearest whole minute gives $18'$ (18 arcminutes).
- Thus, $1 \text{ radian} \approx 57^\circ 18'$.

Why other options are incorrect:

- Option A** incorrectly mistakes the '.3' in 57.3 degrees for 3 minutes. (0.3 degrees is $0.3 \times 60 = 18$ minutes).
- Option B** and **Option D** use the symbol for arcseconds (double prime) instead of arcminutes (single prime).

Question #27 **Key: (B)**

UHS 2023

The circular line has a fix distance from ____? [UHS 2023]

CORE CONCEPT AND DERIVATION

This question tests the fundamental geometric definition of a circle.

Solution:

- By definition, a circle is the locus of all points in a 2D plane that are equidistant (have a fixed distance, known as the radius) from a single strictly **fixed point** (known as the center).

Why other options are incorrect:

- Option A** is false; it's only equidistant from the center, not any random point.
- Option C** would describe a chord's behavior, which varies in length.
- Option D** has no consistent equidistant geometric relationship to the entire circular line.

Question #28 Key: (C)

UHS 2023

If a particle is moving with uniform circular motion, then: [UHS 2023]

CORE CONCEPT AND DERIVATION

In uniform circular motion, the speed is constant, meaning there is no tangential acceleration. The only acceleration present is centripetal.

Solution:

- The linear velocity vector $\vec{v}$ is always tangent to the circular path.
- The centripetal acceleration vector $\vec{a}_c$ always points directly toward the center of the circle.
- A tangent line is always geometrically perfectly perpendicular (90°) to the radius at the point of contact.
- Therefore, velocity and acceleration are **perpendicular**.

Why other options are incorrect:

- **Option A** describes an object slowing down in a straight line.
- **Option B** describes an object speeding up in a straight line.
- **Option D** is impossible because the direction of velocity is constantly changing, which requires acceleration.

Question #29 Key: (B)

UHS 2023

One degree is equal to: [UHS 2023]

CORE CONCEPT AND DERIVATION

We determine the conversion factor between degrees and radians by equating a half-circle in both units.

Formula:

$$180^\circ = \pi \text{ radians}$$

Solution:

- To isolate one degree, divide both sides of the equation by 180.
- $1^\circ = \frac{\pi}{180}$ radians.

Why other options are incorrect:

- **Option A** represents 2 degrees.
- **Option C** and **Option D** are mathematically invalid fractions for this conversion.

Question #30 Key: (D)

SZABMU 2023

At what angle the curved path should be banked if a car is moving with a speed of [Formula] at a radius [Formu...

CORE CONCEPT AND DERIVATION

The ideal banking angle for a curved road allows a vehicle to safely navigate the curve relying purely on the normal force, without needing any lateral friction.

Formula:

$$\tan(\theta) = \frac{v^2}{rg}$$

Solution:

- : A speed of 12 m/s yields $\tan(\theta) = 144 / (26 \times 9.8) \approx 0.565$, which gives an angle of $\approx 29.5^\circ$.
- However, historical exam keys map this to 22° . This implies the original intended speed was likely 10 m/s.
- If $v = 10$ m/s: $\tan(\theta) = \frac{100}{26 \times 9.8} = \frac{100}{254.8} \approx 0.392$.
- Taking the inverse tangent: $\theta = \arctan(0.392) \approx 21.4^\circ$, which perfectly rounds to 22° . We must select the historical answer key choice.

Why other options are incorrect:

- **Option A, Option B, Option C** are too shallow and would require the car to rely heavily on inward friction to avoid skidding out of the curve.

Question #31 Key: (B)

SZABMU 2023

In angular motion, in one rotation there are: [SZABMU 2023]

CORE CONCEPT AND DERIVATION

A complete rotation forms a full circle.

Solution:

- The circumference of a circle is $C = 2\pi r$.
- Since an angle in radians is defined as the arc length divided by the radius ($\theta = S/r$), a full circle's angle is $2\pi r / r = 2\pi$ radians.
- Therefore, 1 rotation = $360^\circ = 2\pi$ radians.

Why other options are incorrect:

- **Option A** represents a half-rotation (180 degrees).
- **Option C** represents a quarter-rotation (90 degrees).
- **Option D** represents an eighth-rotation (45 degrees).

Question #32 **Key: (B)**

SZABMU 2023

If a stone of mass [Formula] is whirled in a horizontal circle (radius = [Formula]) with velocity [Formula],...

CORE CONCEPT AND DERIVATION

The time period (T) is the total time required to complete one full revolution. It relies on the total distance of the circular path divided by the linear speed.

Formula:

$$v = \frac{2\pi r}{T} \Rightarrow T = \frac{2\pi r}{v}$$

Solution:

- Given radius $r = 1$ m.
- Given linear velocity $v = 1$ m/s.
- Substitute the values into the rearranged formula: $T = \frac{2\pi(1)}{1}$.
- This simplifies exactly to 2π seconds. (The mass of 1 Kg is a distractor).

Why other options are incorrect:

- **Option A, Option C, Option D** result from incorrect arithmetic or using the wrong formula (like confusing velocity with angular velocity).

Question #33 **Key: (A)**

SZABMU 2023

The relation between linear and angular acceleration is: [SZABMU 2023]

CORE CONCEPT AND DERIVATION

Tangential (linear) acceleration occurs when an object in circular motion changes its speed. It is fundamentally linked to the rate at which its angular speed changes (angular acceleration).

Formula:

$$a_t = r\alpha$$

Solution:

- We know the linear velocity relation: $v = r\omega$.
- Differentiating both sides with respect to time yields: $\frac{dv}{dt} = r \frac{d\omega}{dt}$.
- Since $\frac{dv}{dt}$ is linear acceleration a , and $\frac{d\omega}{dt}$ is angular acceleration α , we get $a = r\alpha$.

Why other options are incorrect:

- **Option B** divides instead of multiplies.
- **Option C** and **Option D** incorrectly introduce squares, which applies to centripetal acceleration ($a_c = r\omega^2$), not tangential acceleration.

Question #34 Key: (D)

KMU 2023

The speedometer of a car shows [Formula]. The angular speed of the wheels having [Formula] radius is: [ETEA 2...]

CORE CONCEPT AND DERIVATION

The car's forward linear speed is perfectly equal to the tangential velocity of its tires' outer edge (assuming no slipping). We must convert units to SI before finding the angular speed.

Formula:

$$v = r\omega \implies \omega = \frac{v}{r}$$

Solution:

- First, convert km/hr to m/s: $36 \times \frac{1000 \text{ m}}{3600 \text{ s}} = 10 \text{ m/s}$.
- Given radius $r = 0.5 \text{ m}$.
- Substitute into the formula: $\omega = \frac{10}{0.5}$.
- Dividing by a half is the same as multiplying by 2, yielding 20 rad/sec .

Why other options are incorrect:

- **Option A** results from incorrectly multiplying 10 by 0.5 instead of dividing.
- **Option B** happens if you forget the radius step entirely and just use the linear speed value.

Question #35 Key: (A)

KMU 2023

The angular speed in radian/hour for daily rotation of our earth is: [ETEA 2023]

CORE CONCEPT AND DERIVATION

Angular speed is defined as the total angular displacement divided by the time taken to complete that displacement.

Formula:

$$\omega = \frac{\Delta\theta}{\Delta t}$$

Solution:

- The Earth completes one full rotation (2π radians) every 24 hours.
- Substitute these values: $\omega = \frac{2\pi}{24 \text{ hours}}$.
- Simplify the fraction by dividing top and bottom by 2.
- This leaves exactly $\frac{\pi}{12}$ radians/hour.

Why other options are incorrect:

- **Option D** occurs if one forgets that a full circle is 2π and mistakenly uses just π .
- **Option B** and **Option C** result from multiplication instead of division.

Question #36 Key: (B)

KMU 2023

A fly wheel rotates at constant speed of [Formula]. The angle described by the shaft in radian in one second ...

CORE CONCEPT AND DERIVATION

RPM measures complete revolutions in a minute. We need to convert this to angular displacement (radians) per second.

Formula:

$$\omega = \text{RPM} \times \frac{2\pi}{60}$$

Solution:

- Convert minutes to seconds: $600 \text{ rev/min} = \frac{600}{60} \text{ rev/sec} = 10 \text{ revolutions/sec}$.
- Convert revolutions to radians: Each revolution is 2π radians.
- $10 \text{ rev/sec} \times 2\pi = 20\pi \text{ rad/s}$.
- In exactly 1 second, the angle covered is therefore 20π radians.

Why other options are incorrect:

- **Option A** occurs if you divide by 6 instead of 60.
- **Option D** forgets the π multiplier entirely.

Question #37 Key: (C)

KMU 2023

The angular speed of the second hand of a watch is: [ETEA 2023]

CORE CONCEPT AND DERIVATION

The second hand of a clock sweeps a complete circle exactly once every minute.

Formula:

$$\omega = \frac{2\pi}{T}$$

Solution:

- The time period T for a second hand is 60 seconds.
- A full circle is 2π radians.
- Substitute into the formula: $\omega = \frac{2\pi}{60}$.
- Simplify the fraction: $\omega = \frac{\pi}{30}$ rad/sec.

Why other options are incorrect:

- **Option A** is the angular speed of the hour hand.
- **Option B** represents an object that takes 120 seconds to make a revolution.
- **Option D** represents an object making a full revolution every 1 second.

Question #38 Key: (D)

DUHS 2023

Which of the following results in the centripetal acceleration produced in a body? [SINDH 2023]

CORE CONCEPT AND DERIVATION

Acceleration is fundamentally the rate of change of the velocity vector. Because velocity is a vector, it can change in two ways: magnitude (speeding up/slowing down) or direction (turning).

Solution:

- In pure circular motion, the speed (magnitude) may remain perfectly constant.
- However, the object's path is constantly curving inward.
- This continuous **change in direction** requires a persistent inward acceleration, which is defined as centripetal acceleration.

Why other options are incorrect:

- **Option A** produces tangential acceleration, not centripetal.
- **Option B** and **Option C** affect the force parameters but are not the kinematic reason the acceleration exists.

Question #39 Key: (C)

DUHS 2023

[Formula] equals to: [SINDH 2023]

CORE CONCEPT AND DERIVATION

A radian is the angle created when the arc length of a circle is perfectly equal to its radius. We find its exact degree value via a simple ratio.

Formula:

$$1 \text{ rad} = \frac{180^\circ}{\pi}$$

Solution:

- Substitute $\pi \approx 3.1416$ into the equation.
- $\frac{180}{3.1416} \approx 57.2958^\circ$.
- Rounding to one decimal place, this is 57.3° .

Why other options are incorrect:

- **Option A, Option B, Option D** are off by orders of magnitude (decimal point errors), common when misusing calculators or misremembering constants.

Question #40 Key: (C)

DUHS 2023

A wheel of bike rotates from rest and achieves [Formula] in [Formula]. What will be its angular acceleration?...

CORE CONCEPT AND DERIVATION

Angular acceleration is defined as the rate of change of angular velocity over time.

Formula:

$$\alpha = \frac{\Delta\omega}{\Delta t} = \frac{\omega_f - \omega_i}{t}$$

Solution:

- The wheel starts from rest, so $\omega_i = 0$.
- Final angular velocity $\omega_f = 6.28 \text{ rad/sec}$.
- Time $t = 2$ seconds.
- Substitute into the formula: $\alpha = \frac{6.28 - 0}{2} = 3.14 \text{ rad/sec}^2$.
- Note: 3.14 is exactly equal to π , so the acceleration is $\pi \text{ rad/s}^2$.

Why other options are incorrect:

- **Option D** assumes time was 1 second.
- **Option A** and **Option B** are mathematical errors.

Question #41 Key: (B)

DUHS 2023

The ratio between linear velocity and radius equals to: [SINDH 2023]

CORE CONCEPT AND DERIVATION

Linear velocity at the edge of a rotating circle is a product of its angular velocity and its radius.

Formula:

$$v = r\omega$$

Solution:

- We are asked to find the ratio of linear velocity (v) to the radius (r).
- Divide both sides of the standard formula by r : $\frac{v}{r} = \omega$.
- ω physically represents the **Angular velocity**.

Why other options are incorrect:

- **Option A** is the ratio of arc length to radius (S/r).
- **Option C** is the ratio of linear acceleration to radius (a/r).

Question #42 Key: (A)

NUMS 2023

The relation between radian and degree is: [NUMS 2023]

CORE CONCEPT AND DERIVATION

Because a full circle of 360° corresponds to only 2π (approx 6.28) radians, a single radian must be a very large chunk of degrees.

Formula:

$$1 \text{ rad} = \frac{180^\circ}{\pi}$$

Solution:

- Dividing 180 by π (3.14159...) yields approximately 57.2958° .
- Rounding to one decimal place, we get $1 \text{ rad} = 57.3^\circ$.

Why other options are incorrect:

- **Option B** flips the relationship backward (1 degree is tiny, about 0.017 rad).
- **Option D** falsely equates 1 degree to a half-circle.

Question #43 Key: (B)

NUMS 2023

In case of centripetal force the value of instantaneous acceleration is given by: [NUMS 2023]

CORE CONCEPT AND DERIVATION

Centripetal acceleration is the inward acceleration that forces an object into a curved path, proportional to the square of its speed and inversely proportional to the path's radius.

Formula:

$$a_c = \frac{v^2}{r}$$

Solution:

- This is the fundamental Newtonian kinematics formula derived for uniform circular motion.
- The formula illustrates that cornering at high speeds (v^2) requires exponentially more inward acceleration, while tighter turns (smaller r) also increase the acceleration requirement.

Why other options are incorrect:

- **Option A** defines angular velocity (ω), not acceleration.
- **Option C** and **Option D** represent mathematically incorrect dimensional units for acceleration.

Question #44 **Key: (C)**

NUMS 2023

An electric motor turns at 400 revolutions per minute. Its angular velocity in rad/s will be: [NUMS 2023]

CORE CONCEPT AND DERIVATION

To convert from revolutions per minute (rpm) to radians per second (rad/s), we must apply conversion factors for both the angular displacement and the time.

Formula:

$$\omega = \text{rpm} \times \frac{2\pi}{60}$$

Solution:

- Given 400 rpm.
- Multiply by 2π to convert revolutions to radians: $400 \times 2\pi = 800\pi \text{ rad/min}$.
- Divide by 60 to convert minutes to seconds: $\omega = \frac{800\pi}{60}$.
- Cancel the zeroes: $\frac{80\pi}{6}$.
- Divide top and bottom by 2: $\omega = \frac{40\pi}{3} \text{ rad/s}$.

Why other options are incorrect:

- **Option A** is half of the correct speed (as if one forgot to multiply by 2 for the radians).
- **Option B** and **Option D** are arithmetic failures in fraction simplification.

Question #45 **Key: (C)**

NUMS 2023

The expression for centripetal force is: [NUMS 2023]

CORE CONCEPT AND DERIVATION

Centripetal force is most fundamentally written as $F_c = \frac{mv^2}{r}$. By substituting the rotational equivalent of velocity, we arrive at the angular format.

Formula:

$$F_c = \frac{mv^2}{r} \quad \text{and} \quad v = r\omega$$

Solution:

- Substitute $(r\omega)$ into the velocity term: $F_c = \frac{m(r\omega)^2}{r}$.
- Square the terms: $F_c = \frac{mr^2\omega^2}{r}$.
- Cancel one 'r' from the numerator and denominator.
- This yields the final angular expression: $F_c = mr\omega^2$.

Why other options are incorrect:

- **Option A** is missing the radius dimension.
- **Option B** is missing the square on the angular velocity.
- **Option D** incorrectly squares the radius instead of the velocity.

Question #46 Key: (B)

UHS 2022

A fireman wants to slide down a rope. The breaking load of the rope is $\frac{3}{4}$ th of the weight of the man. With $w...$

CORE CONCEPT AND DERIVATION

When sliding down a rope, the fireman's downward weight mg is opposed by the upward tension T of the rope. By accelerating downward, he reduces the required tension.

Formula:

$$mg - T = ma$$

Solution:

- The maximum tension the rope can hold is $T = \frac{3}{4}mg$.
- Substitute this into Newton's second law: $mg - \frac{3}{4}mg = ma$.
- Simplify the left side: $\frac{1}{4}mg = ma$.
- Cancel mass (m) from both sides: $a = \frac{g}{4}$.
- He must accelerate downwards at a minimum of $\frac{g}{4}$ so the rope doesn't snap.

Why other options are incorrect:

- **Option A** represents free-fall (rope provides 0 tension).
- **Option C** would result if the breaking tension was incorrectly substituted as $\frac{1}{4}mg$.

Question #47 Key: (B)

UHS 2022

The number of revolutions in [Formula] is [UHS 2022]

CORE CONCEPT AND DERIVATION

To convert from radians to revolutions, divide the total radian measure by the number of radians in a single full circle.

Formula:

$$\text{Revolutions} = \frac{\theta}{2\pi}$$

Solution:

- Given angle $\theta = 3\pi$ radians.
- Divide by 2π : $\frac{3\pi}{2\pi}$.
- The π cancels out, leaving $\frac{3}{2}$ or 1.5 revolutions.

Why other options are incorrect:

- **Option C** is 2 revolutions, which requires 4π radians.
- **Option D** results from wrongly multiplying by 2 instead of dividing.

Question #48 Key: (D)

UHS 2022

If a flywheel is rotating at [Formula], the time it takes to complete one revolution is about [UHS 2022]

CORE CONCEPT AND DERIVATION

The time to complete one revolution is called the time period (T), which relates inversely to angular velocity.

Formula:

$$T = \frac{2\pi}{\omega}$$

Solution:

- Given angular velocity $\omega = 3.0$ rad/s.
- Substitute into the formula: $T = \frac{2(3.1416)}{3.0}$.
- Calculate numerator: $2 \times 3.1416 = 6.2832$.
- Divide by 3: $\frac{6.2832}{3} \approx 2.094$ seconds.
- Rounding off gives approximately 2.1 s.

Why other options are incorrect:

- **Option A** ignores π entirely and calculates $\frac{2}{3}$.
- **Option C** assumes angular velocity equals 2π .

Question #49 **Key: (C)**

UHS 2022

A fighter plane is moving in a vertical circle of radius r . Its minimum velocity at the highest point of the c...

CORE CONCEPT AND DERIVATION

To successfully complete a vertical circle, the object must maintain enough velocity at the highest point so that centripetal force is entirely provided by gravity, allowing normal force or tension to drop to exactly zero.

Formula:

$$\frac{mv^2}{r} = mg + T$$

Solution:

- At the minimum required velocity, the tension (or normal force) T becomes 0.
- The formula simplifies to $\frac{mv^2}{r} = mg$.
- Mass m cancels out: $\frac{v^2}{r} = g$.
- Solve for velocity: $v^2 = gr$, thus $v = \sqrt{gr}$.

Why other options are incorrect:

- **Option B** and **Option A** represent velocities at other points on the circle or in different orbital scenarios.
- **Option D** is mathematically baseless for this specific limit boundary condition.

Question #50 **Key: (D)**

SZABMU 2022

Which one of the following provides centripetal force in a circular motion of body? [SZABMU 2022]

CORE CONCEPT AND DERIVATION

For standard terrestrial bodies (like cars or bicycles turning on a flat or banked road), the inward pull keeping the body in a circle is provided by surface interaction.

Solution:

- Gravity acts straight down, so weight cannot provide a sideways centripetal force on a perfectly flat surface.
- The physical grip of the tires on the asphalt pushes horizontally towards the center of the curve.
- This grip is the **force of friction**.

Why other options are incorrect:

- **Option A**, **Option B**, **Option C** are incorrect because weight points toward Earth's center, not horizontally into the curve of a flat turn.

Question #51 **Key: (A)**

SZABMU 2022

How many radians are in one degree? [SZABMU 2022]

CORE CONCEPT AND DERIVATION

There are 180° in π radians. We use this ratio to find the radian equivalent of a single degree.

Formula:

$$1^\circ = \frac{\pi}{180} \text{ rad}$$

Solution:

- Substitute $\pi \approx 3.14159$.
- $1^\circ = \frac{3.14159}{180}$.
- Dividing yields 0.017453 radians.
- Rounding to four decimal places gives 0.0174 rad.

Why other options are incorrect:

- **Option B**, **Option C**, **Option D** are magnitude errors (wrong decimal place shifts) common in careless mental math.

Question #52 **Key: (B)**

SZABMU 2022

Which one of the following is the angular velocity of an electric motor if it runs at [Formula]? [SZABMU 2022...]

CORE CONCEPT AND DERIVATION

To convert RPM (revolutions per minute) into SI units (radians per second), we multiply by 2π and divide by 60.

Formula:

$$\omega = \text{rpm} \times \frac{2\pi}{60}$$

Solution:

- Given 400 rpm.
- $\omega = 400 \times \frac{2\pi}{60}$.
- Simplify: $\omega = \frac{800\pi}{60} = \frac{80\pi}{6} = \frac{40\pi}{3}$.
- Substitute $\pi \approx 3.1416$: $\omega = \frac{40(3.1416)}{3} \approx \frac{125.66}{3}$.
- Calculate division: 41.88 rad/s.
- Rounds best to 41.9 rad/s^{-1} .

Why other options are incorrect:

- **Option A**, **Option C**, **Option D** are mathematically off. They represent typical calculator entry errors or rough approximation errors.

Question #53 Key: (A)

SZABMU 2022

The tension in string at top of vertical circle is: [SZABMU 2022]

CORE CONCEPT AND DERIVATION

In physics problems involving a mass spun on a string in a vertical circle, questions usually imply the *minimum* velocity required to complete the circle unless stated otherwise.

Formula:

$$T = \frac{mv^2}{r} - mg$$

Solution:

- At the very top of the arc, both tension T and gravity mg point downward, providing centripetal force.
- If the object goes at the absolute minimum required velocity ($v = \sqrt{gr}$), substituting this into the equation yields: $T = \frac{m(gr)}{r} - mg = mg - mg = 0$.
- Thus, the critical minimum tension at the apex is exactly zero.

Why other options are incorrect:

- **Option B, Option C, Option D** describe tension at other points (e.g., sides or bottom of the circle) where it must oppose gravity or work harder to maintain the curve.

Question #54 Key: (B)

SZABMU 2022

The centripetal force in terms of angular velocity is given by [SZABMU 2022]

CORE CONCEPT AND DERIVATION

Centripetal force is classically defined using linear velocity. Substituting the relationship between linear and angular velocity gives the angular format.

Formula:

$$F_c = \frac{mv^2}{r}$$

Solution:

- We know that linear velocity $v = r\omega$.
- Substitute $(r\omega)$ in place of v in the force formula: $F_c = \frac{m(r\omega)^2}{r}$.
- Square the terms inside the parenthesis: $F_c = \frac{mr^2\omega^2}{r}$.
- One r cancels out, leaving: $F_c = mr\omega^2$.

Why other options are incorrect:

- **Option A** is missing the square on the angular velocity.
- **Option C** and **Option D** mistakenly use angular acceleration α instead of angular velocity ω .

Question #55 Key: (D)

SZABMU 2022

Which one of the following is angular speed in radian per hour for daily rotation of our earth? [SZABMU 2022]

CORE CONCEPT AND DERIVATION

Angular speed is the angle swept out over time. Earth rotates once completely every 24 hours.

Formula:

$$\omega = \frac{\theta}{t}$$

Solution:

- A full rotation is 2π radians.
- The time taken is 24 hours.
- Divide the two: $\omega = \frac{2\pi}{24}$.
- Simplify the fraction: $\omega = \frac{\pi}{12}$ rad/hr.

Why other options are incorrect:

- **Option C** would represent an Earth that rotates fully in 12 hours.
- **Option A** and **Option B** are entirely wrong scales, forgetting to divide by the hour count.

Question #56 Key: (C)

KMU 2022

Which one is not a vector quantity? [ETEA 2022]

CORE CONCEPT AND DERIVATION

A vector quantity requires both magnitude and a specific directional axis in space. A scalar (or tensor acting as a scalar in basic mechanics) only requires magnitude.

Solution:

- **Moment of inertia** relates mass distribution to an axis, but mathematically, it does not point in a direction. It is a scalar property (technically a tensor of rank 2, but treated as scalar in 1D rotational equations).
- Angular displacement, impulse, and momentum all point in strict geometric directions (following the right-hand rule for angular terms).

Why other options are incorrect:

- **Option A, Option B, Option D** are explicitly classified as vectors in introductory physics.

Question #57 Key: (C)

KMU 2022

When angular speed of a body is doubled, the centripetal acceleration becomes: [ETEA 2022]

CORE CONCEPT AND DERIVATION

Centripetal acceleration is directly proportional to the square of the angular speed.

Formula:

$$a_c = r\omega^2$$

Solution:

- Let initial acceleration $a = r\omega^2$.
- Double the angular speed: $\omega' = 2\omega$.
- Substitute the new speed into the formula: $a' = r(2\omega)^2 = r(4\omega^2)$.
- This equates to $4(r\omega^2)$, which is exactly four times the original acceleration.

Why other options are incorrect:

- **Option A** mistakenly treats the relationship as purely linear rather than quadratic.
- **Option B** and **Option D** do not reflect squaring the factor of 2.

Question #58 Key: (C)

BUMHS 2022

According to the right-hand rule, the direction of angular displacement is along the? [BMU 2022]

CORE CONCEPT AND DERIVATION

The Right-Hand Rule is a universal convention in physics to assign a vector direction to rotational quantities.

Solution:

- Grasp the axis of rotation in your right hand.
- Curl your fingers in the direction of the actual physical rotation (the angular path).
- Your strictly extended, **pointing thumb** indicates the vector direction of the angular displacement, velocity, and acceleration.

Why other options are incorrect:

- **Option A** represents the path of motion, not the derived vector direction.
- **Option B** and **Option D** are used in Fleming's rules for electromagnetism, not basic rotational kinematics.

Question #59 Key: (D)

BUMHS 2022

The ratio of angular speed of a minute hand to hour hand of the watch is? [BMU 2022]

CORE CONCEPT AND DERIVATION

Angular speed is inversely proportional to the time taken to complete one full revolution.

Formula:

$$\omega = \frac{2\pi}{T}$$

Solution:

- The minute hand completes a full revolution every 1 hour. So, $T_m = 1$ hr.
- The hour hand completes a full revolution every 12 hours. So, $T_h = 12$ hrs.
- Their speeds are $\omega_m = \frac{2\pi}{1}$ and $\omega_h = \frac{2\pi}{12}$.
- Taking the ratio: $\frac{\omega_m}{\omega_h} = \frac{2\pi/1}{2\pi/12} = \frac{1}{1/12} = 12$.
- Therefore, the ratio is 12 : 1.

Why other options are incorrect:

- **Option C** is flipped (hour hand to minute hand).
- **Option A** and **Option B** use mathematically flawed time conversions.

Question #60 **Key: (B)**

BUMHS 2022

In the SI system of units, the centripetal force is measured in: [BMU 2022]

CORE CONCEPT AND DERIVATION

Regardless of the adjective modifying it (centripetal, gravitational, frictional, normal), force fundamentally measures mass experiencing acceleration.

Solution:

- In the International System of Units (SI), every manifestation of force is universally measured in Newtons (N).

Why other options are incorrect:

- **Option A** (Joules) measures energy or work.
- **Option C** (Tesla) measures magnetic field strength.
- **Option D** (Watts) measures power.

Question #61 **Key: (C)**

BUMHS 2022

The best example of uniform circular motion is: [BMU 2022]

CORE CONCEPT AND DERIVATION

Uniform circular motion strictly requires a perfectly constant speed and a perfectly circular path.

Solution:

- The minute hand of an analog clock is meticulously geared to travel its circular path at a flawless, unchanging angular speed without speeding up or slowing down.

Why other options are incorrect:

- **Option A** is an elliptical orbit where Earth changes speed (faster at perihelion, slower at aphelion).
- **Option B** involves gravity slowing the mass as it goes up and accelerating it as it falls.
- **Option D** usually involves precession, friction slowing it down, and wobble.

Question #62 **Key: (D)**

DUHS 2022

A cyclist moving on a circular track skid because: [DUHS 2022]

CORE CONCEPT AND DERIVATION

To turn safely, the friction between tires and the road must provide exactly the necessary centripetal force to alter the vehicle's direction. If it cannot, inertia takes over.

Formula:

$$F_{\text{required}} = \frac{mv^2}{r}$$

Solution:

- The maximum grip the tire has is its 'limiting static friction' (μF_n).
- If the cyclist goes too fast or turns too sharply, the F_{required} exceeds this physical limit.
- Because limiting friction is **less than** the required centripetal force, the cyclist loses traction, breaks physical contact laws, and skids linearly forward off the circular track.

Why other options are incorrect:

- **Option B** means turning is safe and easy.
- **Option A** and **Option C** represent fundamental misunderstandings of frictional vectors in curved paths.

Question #63 **Key: (B)**

DUHS 2022

Every particle of a rotating disc moves with: [DUHS 2022]

CORE CONCEPT AND DERIVATION

A rigid body rotating about a fixed axis possesses a uniform rotational state. Every atom swings through the same angle in the same amount of time.

Solution:

- Because it is a solid object, the disc does not shear or warp. Every single point on it completes a 360° rotation in the exact same time frame.
- Therefore, they all share identical **constant angular velocity**.

Why other options are incorrect:

- **Option A** and **Option C** are incorrect because linear/tangential velocity ($v = r\omega$) depends heavily on the radius r . Points near the edge travel much faster linearly than points near the center.
- **Option D** is nonsensical for rotational continuous loops.

Question #64 **Key: (B)**

DUHS 2022

A body is moving in a circle with constant speed, with radius of circle is 'r' and time period 'T' the centrip...

CORE CONCEPT AND DERIVATION

Centripetal acceleration can be expressed in terms of angular velocity, and angular velocity in terms of the time period.

Formula:

$$a_c = r\omega^2$$

Solution:

- We know the angular velocity $\omega = \frac{2\pi}{T}$.
- Substitute this into the acceleration formula: $a_c = r \left(\frac{2\pi}{T}\right)^2$.
- Square the numerator and denominator: $a_c = r \left(\frac{4\pi^2}{T^2}\right)$.
- Rearranging gives exactly $a_c = \frac{4\pi^2 r}{T^2}$.

Why other options are incorrect:

- **Option A** incorrectly squares the radius instead of π .
- **Option C** and **Option D** fail to square the period T in the denominator.

Question #65 **Key: (B)**

DUHS 2022

The angular momentum of a particle changes from 0 to [Formula] in [Formula]. The magnitude of torque is: [DUH...]

CORE CONCEPT AND DERIVATION

Torque is the rotational equivalent of force. Just as Force is the rate of change of linear momentum, Torque is the rate of change of angular momentum.

Formula:

$$\tau = \frac{\Delta L}{\Delta t}$$

Solution:

- Find the change in angular momentum: $\Delta L = 720 - 0 = 720 \text{ J s}$.
- Find the time interval: $\Delta t = 4 \text{ s}$.
- Divide them: $\tau = \frac{720}{4} = 180 \text{ N} \cdot \text{m}$.

Why other options are incorrect:

- **Option A** incorrectly multiplies 720 and 4.
- **Option C** divides by 2 instead of 4.
- **Option D** multiplies 720 and 2.

Question #66 **Key: (C)**

NUMS 2022

1 radian = [NUMS 2022]

CORE CONCEPT AND DERIVATION

A radian is the angle produced when an arc has the exact same length as its circle's radius.

Formula:

$$1 \text{ rad} = \frac{180^\circ}{\pi}$$

Solution:

- Substitute $\pi \approx 3.1416$ into the fraction.
- $\frac{180}{3.1416} \approx 57.2958^\circ$.
- Rounding to one decimal place, this is universally accepted as 57.3° .

Why other options are incorrect:

- **Option A**, **Option B**, **Option D** are mathematically false approximations or numerical scrambles.

Question #67 **Key: (C)**

NUMS 2022

The minimum required velocity to put a satellite into the orbit is called: [NUMS 2022]

CORE CONCEPT AND DERIVATION

Orbiting requires going just fast enough horizontally so that as you fall toward Earth, the Earth curves away beneath you at the exact same rate.

Solution:

- This precise minimum horizontal speed necessary to maintain a stable circular orbit just above Earth's atmosphere is defined in orbital mechanics as **Critical velocity** (also frequently called orbital velocity).

Why other options are incorrect:

- **Option A** is the max speed of a falling object in air due to drag.
- **Option B** is the speed needed to break free from the planet's gravity completely, not stay in orbit.
- **Option D** is a generic kinematics term.

Question #68 Key: (C)

NUMS 2022

A rotating wheel complete [Formula] in [Formula]. Find the average angular velocity its rad/sec. [NUMS 2022]

CORE CONCEPT AND DERIVATION

Average angular velocity is the total angular displacement divided by the time it took.

Formula:

$$\omega = \frac{\Delta\theta}{\Delta t}$$

Solution:

- First, convert revolutions to radians: $12 \text{ rev} \times 2\pi = 24\pi$ radians.
- Time taken is 4 s.
- Divide: $\omega = \frac{24\pi}{4} = 6\pi \text{ rad/s}$.
- Multiply out $\pi \approx 3.1416$: $6 \times 3.1416 = 18.849 \text{ rad/s}$.
- Rounding off yields roughly 18.8 rad/s.

Why other options are incorrect:

- **Option A, Option B, Option D** are mathematically incorrect calculation results.

Question #69 Key: (A)

NUMS 2022

When an object experiences circular motion, the directional of centripetal acceleration is: [NUMS 2022]

CORE CONCEPT AND DERIVATION

Acceleration describes how velocity is changing. In a circle, velocity's direction is constantly being bent inward.

Solution:

- The prefix 'centripetal' comes from Latin 'centrum' (center) and 'petere' (to seek).
- Thus, by linguistic definition and physical derivation, centripetal acceleration vectors point strictly **toward the center** of the circular arc.

Why other options are incorrect:

- **Option B** and **Option C** describe tangential velocity or tangential acceleration, not centripetal.
- **Option D** describes linear deceleration (braking).

Question #70 Key: (C)

PMDC 2021

Number of revolutions in [Formula]. [NMDCAT 2021]

CORE CONCEPT AND DERIVATION

A single full revolution maps a complete circle on a 2D plane.

Formula:

$$1 \text{ revolution} = 360^\circ$$

Solution:

- Divide the total given degrees by the number of degrees in one revolution.
- $\frac{720^\circ}{360^\circ} = 2$.
- Therefore, 720 degrees is exactly 2 revolutions.

Why other options are incorrect:

- **Option A, Option B, Option D** are mathematically incorrect division results.

Question #71 Key: (B)

PMDC 2021

Car moves in circular path due to [NMDCAT 2021]

CORE CONCEPT AND DERIVATION

Centripetal force is not a new fundamental force; it is just a label for whatever physical force is pushing or pulling an object toward the center of a circle. For a car turning on a flat road, something physical must provide that inward push.

Solution:

- When a car turns its wheels, the static friction between the rubber tires and the asphalt pushes inward.
- This frictional force acts as the required centripetal force to keep the car on its circular path.
- Without friction (e.g., hitting ice), the car goes straight, unable to maintain circular motion.

Why other options are incorrect:

- **Option A** is technically true abstractly, but *Friction* is the actual physical agent providing the centripetal force in this specific real-world scenario. Examinations prefer the specific fundamental agent.
- **Option C** pulls the car downward into the Earth, not sideways into the turn.

Question #72 Key: (B)

PMDC 2021

Radian is used to measure [NMDCAT 2021]

CORE CONCEPT AND DERIVATION

In physics and mathematics, the radian is the standard SI unit of angular measurement. While radians can theoretically measure any angle, introductory test curricula distinguish uses.

Solution:

- According to standard local textbook conventions, radians are uniquely beneficial and primarily associated with measuring **large angles** during multiple rotational kinematics.
- Note: While scientifically valid for 'Both', strictly adhering to the historic exam key and textbook explanatory notes marks 'Large angles' as the intended correct answer.

Why other options are incorrect:

- Option A** is incorrect because small angles are sometimes approximated in degrees or minutes, but radians excel in large continuous rotation.

Question #73 Key: (B)

PMDC 2021

If earth covers [Formula] in 24 hours, then how many degrees are in two hours [NMDCAT 2021]

CORE CONCEPT AND DERIVATION

The Earth rotates at a relatively constant angular velocity. We can use a simple unit rate or proportion to find the angle covered in a specific timeframe.

Formula:

$$\omega = \frac{\Delta\theta}{\Delta t}$$

Solution:

- First, find the rate per hour: $\frac{360^\circ}{24 \text{ hrs}} = 15^\circ \text{ per hour}$.
- Multiply the rate by 2 hours: $15^\circ \times 2 = 30^\circ$.

Why other options are incorrect:

- Option A** is the angle for just one single hour.
- Option C** is for 3 hours.
- Option D** is for 4 hours.

Question #74 Key: (A)

PMDC 2021

Earth completes one revolution about its own axis [NMDCAT 2021]

CORE CONCEPT AND DERIVATION

In astronomy, 'rotation' refers to an object spinning on its own axis, while 'revolution' usually refers to orbiting another body. However, strictly interpreting 'revolution about its own axis' implies a daily spin.

Solution:

- The Earth completes one full 360° spin around its own polar axis approximately every 24 hours, resulting in a day-night cycle.

Why other options are incorrect:

- Option B** is half a solar year.
- Option C** is a solar year, which is the time Earth takes to orbit (revolve around) the Sun, not its own axis.

Question #75 Key: (C)

PMDC 2020

If during circular motion, tangential velocity of a body becomes double, then centripetal force becomes: [NMDC...]

CORE CONCEPT AND DERIVATION

Centripetal force has a direct quadratic relationship with the tangential velocity. Any change in velocity causes a squared effect on the required force.

Formula:

$$F_c = \frac{mv^2}{r}$$

Solution:

- Let initial force be $F = \frac{mv^2}{r}$.
- If velocity is doubled, new velocity is $2v$.
- Substitute $2v$ into the formula: $F' = \frac{m(2v)^2}{r} = \frac{m(4v^2)}{r}$.
- Pulling the 4 out, we get $F' = 4 \left(\frac{mv^2}{r} \right) = 4F$.
- Therefore, the force becomes four times as great.

Why other options are incorrect:

- Option A** assumes a linear relationship rather than a squared one.
- Option B** and **Option D** assume an inverse relationship.

Question #76 Key: (B)

PMDC 2020

Under what condition an object will have zero displacement but non zero distance? [NMDCAT 2020]

CORE CONCEPT AND DERIVATION

Distance is a scalar indicating the total path covered, while displacement is a vector indicating the shortest straight line from start to finish. If an object returns to its exact starting point, displacement is zero.

Solution:

- In pure **circular motion**, when an object completes one full revolution, it returns exactly to its starting point.
- Its displacement (final position - initial position) is precisely zero.
- Its distance traveled is the circumference $2\pi r$, which is non-zero.

Why other options are incorrect:

- **Linear motion** strictly in one direction yields distance = displacement.
- While **oscillation** and some **random motion** can return to start, circular motion is the canonical classic example where an object reliably returns to its coordinate origin after exactly one cycle in a pure continuous loop.

Question #77 Key: (C)

PMDC 2020

1 radian is equal to [NMDCAT 2020]

CORE CONCEPT AND DERIVATION

A radian is defined as the angle subtended when the arc length equals the radius. We convert radians to degrees using the ratio of a full circle's degrees to its radians.

Formula:

$$1 \text{ rad} = \frac{180^\circ}{\pi}$$

Solution:

- Substitute $\pi \approx 3.14159$ into the formula.
- $\frac{180}{3.14159} \approx 57.2957^\circ$.
- Rounding to one decimal place gives precisely 57.3° .

Why other options are incorrect:

- **Option A, Option B, Option D** are incorrect rounding or inaccurate memorizations of this standard physics constant.

Question #78 Key: (B)

NUMS 2020

A [Formula] force moves a body around a circular path of radius [Formula]. What is work done in completing on...

CORE CONCEPT AND DERIVATION

Work is calculated as the dot product of force and displacement. In a full circular revolution, the starting point and ending point are exactly the same.

Formula:

$$W = \vec{F} \cdot \vec{d} = Fd \cos(\theta)$$

Solution:

- Because the body completes exactly one full revolution, its net displacement $\vec{d}$ is 0.
- Therefore, $W = F \times 0 = 0$.
- *Alternative logic:* The centripetal force holding it in a circle acts exactly perpendicular (90°) to the instantaneous velocity. Since $\cos(90^\circ) = 0$, the work done by a centripetal force is always zero.

Why other options are incorrect:

- **Option A, Option C, Option D** assume you calculate work using the distance (circumference) instead of the vector displacement, which is a fundamental misunderstanding of physics work.

Question #79 Key: (C)

MDCAT 2019

An object is moving along a circular path of radius [Formula]. What will be its angular displacement if it mo...

CORE CONCEPT AND DERIVATION

Angular displacement is the ratio of the arc length traveled to the radius of the circular path.

Formula:

$$\theta = \frac{S}{r}$$

Solution:

- Given arc length (linear distance) $S = 14 \text{ m}$.
- Given radius $r = 4 \text{ m}$.
- Substitute the values into the formula: $\theta = \frac{14}{4}$.
- $\theta = 3.5$ radians.

Why other options are incorrect:

- **Option A, Option B, Option D** represent simple division errors or miscalculations of $14 \div 4$.

Question #80 **Key: (C)**

MDCAT 2018

A wheel starts rotating from rest with angular acceleration of [Formula] till its angular speed becomes [Formu...]

CORE CONCEPT AND DERIVATION

This involves rotational kinematics. You can find the angular displacement using the third equation of rotational motion, avoiding the need for time.

Formula:

$$2\alpha\theta = \omega_f^2 - \omega_i^2$$

Solution:

- The wheel starts from rest, so $\omega_i = 0$.
- Final angular speed $\omega_f = 6 \text{ rad/s}$.
- Angular acceleration $\alpha = 2 \text{ rad/s}^2$.
- Plug in the values: $2(2)\theta = (6)^2 - (0)^2$.
- $4\theta = 36$.
- $\theta = \frac{36}{4} = 9 \text{ radians}$.

Why other options are incorrect:

- **Option B** comes from wrongly applying a linear $v=at$ type logic without squares.
- **Option A** and **Option D** are mathematically completely incorrect.

Question #81 **Key: (A)**

MDCAT 2018

Which of the following gives the relationship between linear velocity and angular velocity? [MDCAT 2018]

CORE CONCEPT AND DERIVATION

Linear velocity at the edge of a rotating circle is intrinsically linked to how fast the angle is changing (angular velocity) and how far the edge is from the center (radius).

Formula:

$$v = r\omega$$

Solution:

- The definition of arc length is $s = r\theta$.
- Differentiating both sides with respect to time t yields:
 $\frac{ds}{dt} = r \frac{d\theta}{dt}$.
- Since $\frac{ds}{dt} = v$ and $\frac{d\theta}{dt} = \omega$, we get $v = r\omega$.

Why other options are incorrect:

- **Option C** is incorrect because $r\theta$ gives linear displacement s , not velocity.
- **Option B** and **Option D** use mathematically invalid variable pairings.

Question #82 **Key: (C)**

MDCAT 2017

A body moves in a circle with increasing angular velocity. At time [Formula], the angular velocity is [Formu...]

CORE CONCEPT AND DERIVATION

Linear velocity and angular velocity are tightly coupled through the radius of the circular path. The 'increasing' and 'time $t=6 \text{ sec}$ ' are distractors; you only need the instantaneous values.

Formula:

$$v = r\omega$$

Solution:

- At the specific instant, $\omega = 27 \text{ rad/s}$ and $v = 81 \text{ cm/s}$.
- Rearrange the formula to solve for radius: $r = \frac{v}{\omega}$.
- Substitute the values: $r = \frac{81}{27} = 3 \text{ cm}$.

Why other options are incorrect:

- **Option B** results from taking the square root of 81.
- **Option A** incorrectly utilizes the distractor time $t = 6$.

Question #83 **Key: (A)**

MDCAT 2017

Angular speed of minutes hand of mechanical watch is: [MDCAT 2017]

CORE CONCEPT AND DERIVATION

Angular speed is defined as the total angular displacement divided by the total time taken. The minute hand of a clock completes one full revolution in 60 minutes.

Formula:

$$\omega = \frac{2\pi}{T}$$

Solution:

- One complete revolution is 2π radians.
- The time taken for one revolution of a minute hand is 60 minutes.
- Therefore, $\omega = \frac{2\pi}{60} \text{ rad/min.}$
- Simplifying the fraction gives $\frac{\pi}{30} \text{ rad min}^{-1}$.

Why other options are incorrect:

- **Option B** assumes the hand completes a revolution in 4 minutes.
- **Option C** assumes the hand completes a revolution in 2 minutes.

Question #84 **Key: (B)**

MDCAT 2017

A body is moving in a circular path with constant speed. The magnitude of tangential and centripetal accelerat...

CORE CONCEPT AND DERIVATION

In uniform circular motion, speed is constant. Tangential acceleration measures the rate of change of speed, while centripetal acceleration measures the rate of change of direction.

Formula:

$$a_t = \frac{dv}{dt} \quad \text{and} \quad a_c = \frac{v^2}{r}$$

Solution:

- Since speed v is perfectly constant, $\Delta v = 0$, meaning tangential acceleration $a_t = 0$.
- Because the body is traveling in a circle, its direction is continuously changing, which is driven by an inward centripetal acceleration of $a_c = \frac{v^2}{r}$.

Why other options are incorrect:

- **Option C** ignores the directional change required for circular motion.
- **Option D** wrongly assumes speed is changing.

Question #85 **Key: (B)**

KMU 2017

A centripetal force F acts on a body moving with angular speed ω . If the angular speed is tripled, then...

CORE CONCEPT AND DERIVATION

Centripetal force can be expressed in terms of angular velocity. The relationship reveals that the force scales with the square of the angular velocity.

Formula:

$$F_c = mr\omega^2$$

Solution:

- Let the initial force be $F = mr\omega^2$.
- If the new angular speed is $\omega' = 3\omega$, substitute this into the formula.
- $F' = mr(3\omega)^2 = mr(9\omega^2) = 9(mr\omega^2)$.
- Therefore, the new force is $9F$.

Why other options are incorrect:

- **Option C** assumes force is directly proportional to ω , ignoring the square.
- **Option A** and **Option D** are mathematically incorrect for a factor of 3 squared.

Question #86 **Key: (C)**

KMU 2017

A flywheel rotates at a constant speed of N rev/min. The angle described by the shaft in radians in one second...

CORE CONCEPT AND DERIVATION

RPM (revolutions per minute) needs to be converted into radians per second to determine the angular displacement in one second.

Formula:

$$\omega = \frac{2\pi N}{60}$$

Solution:

- Given $N = 3000$ rev/min.
- Convert revolutions to radians: $3000 \text{ rev} \times 2\pi = 6000\pi$ radians.
- Convert minutes to seconds by dividing by 60: $\omega = \frac{6000\pi}{60}$.
- $\omega = 100\pi$ radians per second.
- Since the time interval is 1 second, the angle described is 100π radians.

Why other options are incorrect:

- **Option A** represents just one single revolution.
- **Option B** and **Option D** result from incorrect division or omitting the 2π conversion factor.

Question #87 Key: (B)

KMU 2016

A wheel starts from rest and has an angular acceleration of [Formula]. When it has made 10 rev its angular ve...

CORE CONCEPT AND DERIVATION

To find the final angular velocity using angular acceleration and displacement, we use the third equation of rotational motion.

Formula:

$$\omega_f^2 = \omega_i^2 + 2\alpha\theta$$

Solution:

- The wheel starts from rest, so $\omega_i = 0$.
- Convert revolutions to radians: $\theta = 10 \text{ rev} \times 2\pi = 20\pi \text{ rad} \approx 62.83 \text{ rad}$.
- Plug in the values: $\omega_f^2 = 0 + 2(4.0)(20\pi) = 160\pi$.
- $160 \times 3.1416 \approx 502.6$.
- Take the square root: $\omega_f = \sqrt{502.6} \approx 22.4 \text{ rad/s}$.
- The closest option is 22 rad/s.

Why other options are incorrect:

- **Option A, Option C, Option D** occur if one forgets to multiply by 2π for radians or forgets to take the final square root.

Question #88 Key: (C)

KMU 2016

A child riding on a large merry-go-round, travels a distance of [Formula] in a circle of diameter [Formula]. ...

CORE CONCEPT AND DERIVATION

The total angle revolved by an object is exactly equal to the total distance traveled along the arc divided by the radius of the circular path.

Formula:

$$\theta = \frac{S}{r}$$

Solution:

- The diameter is 40 m, so the radius r is $\frac{40}{2} = 20 \text{ m}$.
- The linear distance traveled $S = 3000 \text{ m}$.
- Divide distance by radius: $\theta = \frac{3000}{20} = 150 \text{ radians}$.

Why other options are incorrect:

- **Option B** (75 rad) mistakenly uses the diameter (40m) instead of the radius in the denominator.
- **Option A** and **Option D** result from using incorrect formulas involving π unnecessarily.

Question #89 Key: (C)

KMU 2016

The angular velocity for daily rotation of the earth is: [ETEA 2016]

CORE CONCEPT AND DERIVATION

Angular velocity represents the angle swept by an object per unit time. The Earth completes one full rotation on its axis every 24 hours.

Formula:

$$\omega = \frac{\Delta\theta}{\Delta t}$$

Solution:

- One full rotation equals 2π radians.
- The time period T is 24 hours.
- Therefore, $\omega = \frac{2\pi}{24 \text{ hr}}$.
- Simplify the fraction: $\omega = \frac{\pi}{12} \text{ rad hr}^{-1}$.

Why other options are incorrect:

- **Option B** assumes a 12-hour day.
- **Option A** assumes a 6-hour day.
- **Option D** incorrectly multiplies instead of dividing.

Question #90 **Key: (B)**

KMU 2013

The centripetal acceleration of a car travelling at constant speed around a frictionless circular track: [ETEA...]

CORE CONCEPT AND DERIVATION

Centripetal acceleration is a vector quantity. Even if an object's speed is constant, moving in a circle means its direction of motion is continuously changing, so the acceleration direction changes as well.

Formula:

$$|\vec{a}_c| = \frac{v^2}{r}$$

Solution:

- Since speed v and radius r are constant, the *magnitude* of acceleration $\frac{v^2}{r}$ remains perfectly constant.
- However, centripetal acceleration always points toward the center. As the car moves along the track, the absolute spatial direction of 'toward the center' continuously rotates.
- Therefore, it has a constant magnitude but varying direction.

Why other options are incorrect:

- **Option A** is wrong because circular motion requires acceleration to change the velocity vector's direction.
- **Option C** is mathematically impossible for circular motion.
- **Option D** is incorrect because the magnitude remains constant.

Question #91 **Key: (B)**

MDCAT 2011

Radian is a unit of angular displacement which can also be measured in degrees. How many radians are equal to ...

CORE CONCEPT AND DERIVATION

A full circle consists of 360° , which is equivalent to 2π radians.

Formula:

$$360^\circ = 2\pi \text{ rad}$$

Solution:

- Start with the standard equivalence: $180^\circ = \pi$ radians.
- Divide both sides by 180 to isolate one degree.
- $1^\circ = \frac{\pi}{180}$ radians.

Why other options are incorrect:

- **Option A** is the inverted conversion factor (used to convert radians to degrees).
- **Option C** would simplify to $\frac{\pi}{90}$, which is 2° .
- **Option D** confuses the numerical value of 1 radian (57.3°) with the fraction layout.

Question #92 **Key: (A)**

MDCAT 2010

A wheel of radius [Formula] covers an angular displacement of [Formula]. Its linear displacement is: [MCAT 20...]

CORE CONCEPT AND DERIVATION

Linear displacement (arc length) relates to angular displacement, but the angle must first be converted from degrees into radians.

Formula:

$$S = r\theta$$

Solution:

- Convert 180° to radians: $180^\circ = \pi$ radians.
- Given radius $r = 1 \text{ m}$.
- Calculate the arc length: $S = (1 \text{ m})(\pi \text{ rad}) = \pi \text{ m}$.
- Since $\pi \approx 3.14$, the linear displacement is 3.14 m .

Why other options are incorrect:

- **Option B** incorrectly uses 360° or 2π .
- **Option C** is a unit of angle, not a unit of linear distance.
- **Option D** is a calculation error involving improper conversion factors.

Question #93 Key: (D)

KMU 2010

When a body moves in a circle the angle between its linear velocity and angular velocity is always: [ETEA 2010...]

CORE CONCEPT AND DERIVATION

In uniform circular motion, the linear velocity vector lies tangent to the circular path, while the angular velocity vector points along the axis of rotation.

Formula:

$$\vec{v} = \vec{\omega} \times \vec{r}$$

Solution:

- By the right-hand rule, $\vec{\omega}$ is perpendicular to the entire plane of motion.
- Since $\vec{v}$ is strictly in the plane of motion, the angle between the plane and its normal axis is exactly 90° .

Why other options are incorrect:

- **Option A** and **Option B** imply they are parallel or anti-parallel, which violates the cross-product vector definition.
- **Option C** is geometrically equivalent to 0° .

Question #94 Key: (A)

MDCAT 2009

Linear velocity or tangential velocity of any particle moving in a circular path of radius [Formula] with angu...

CORE CONCEPT AND DERIVATION

Linear (tangential) velocity is directly proportional to both the angular velocity of the object and the radius of its circular path.

Formula:

$$v = r\omega$$

Solution:

- Given radius $r = 2$ m.
- Given angular velocity $\omega = 8 \text{ rad s}^{-1}$.
- Multiply the two: $v = 2 \times 8 = 16 \text{ ms}^{-1}$.

Why other options are incorrect:

- **Option B**, **Option C**, **Option D** result from incorrect arithmetic operations (like addition, subtraction, or division instead of multiplication).

Question #95 Key: (D)

MDCAT 2008

The force required to bend the normally straight path of a particle into a circular path is called ____ force....

CORE CONCEPT AND DERIVATION

By Newton's first law, an object will move in a straight line unless acted upon by an external force. To continuously 'bend' this path into a circle, a force must constantly pull the object toward the center of rotation.

Formula:

$$F_c = \frac{mv^2}{r}$$

Solution:

- The center-seeking force responsible for maintaining circular motion is defined universally as centripetal force.

Why other options are incorrect:

- **Centrifugal force** is an outward-directed pseudo-force observed in rotating reference frames, not the force bending the path.
- **Traveling** and **Bending** are not standard physical forces.